

## STUDY ON THE ANTIBACTERIAN CHARACTERISTICS OF TEA TREE OIL AND ITS APPLICATION IN COSMETICS

Phan Dinh Tuan, Hoang Minh Nam, Nguyen Thi Thanh Nga  
University of Technology, VNU-HCM

### 1. INTRODUCTION

Tea trees have been widely planted in the Mekong River Delta area of Vietnam. The investigation has pointed out that its essential oil composition is of about 2% by weight. The essential oil could be easily recovered and purified by steam distillation followed by fractional one [9].

It has been reported that tea tree oil has rather high antibacterial activity [9,12,14,17,18] for its high composition of terpinen-4-ol. Nevertheless, it has been found that not only terpinen-4-ol but also other components have the activity. It is necessary to find out the available minimum dose of the essential oil in practical use that the antibacterial activity could be retained. Besides, it is important to point out the available ability to use different fractions of the essential oil for antibacterial purpose.

### 2. MATERIALS AND METHODS

#### 2.1. Materials

The original tea tree essential oil is recovered and fractionally distilled from the tea trees planted in the Mekong River Delta area of Vietnam. From these fractions, initial essential oil has been prepared by simple mixing the fractions of 0% and 98% terpinen-4-ol to receive the mixtures of about 25%, 50%, 75% and 90% terpinen-4-ol.

#### 2.2. Microorganisms

*Escherichia coli* and *Staphylococcus aureus* microorganisms are supplied by the Department of Biotechnology of Hochiminh City University of Technology. *Pseudomonas aeruginosa* and *Candida albicans* are supplied by the Pasteur Institute at Hochiminh City.

#### 2.3. Methodology

Antibacterial activity of the essential oil, its distilled fractions or of essential oil based cosmetics are determined by the method of diffusion well (hole boring) and the method of microorganism counting. In the case of cosmetic products, the method of microorganism counting is applied, provided that the water-undissolved essential oil is emulsified by Tween 80 as the emulsifying reagent.

### 3. RESULTS AND DISCUSSION

#### 3.1. Qualitative investigation of antibacterial activity by diffusion well method

The hole boring method is applied to all types of chosen microorganisms. The concentrations of the used essential oil fractions are represented by the number of microliters applied to the hole. The investigated results are shown on Figures 1a, 1b, 1c, 1d for *Escherichia coli*, *Staphylococcus aureus*, *Candida albicans* and *Pseudomonas aeruginosa*, respectively.

Checking the effect of essential oil evaporation by contacting the agar medium with the essential oil vapor is shown in Table 1. It has pointed out that only the diffusion of the oil determines the antibacterial characteristics.

It is pointed out from the experimental results that at lower concentration of essential oil, diameters of the antibacterial cycles rapidly increase with its used amounts. At higher concentration of essential oil, the diameters increase also but more slowly. It has the reason of

non-polar nature of the essential oil, so that its diffusion in polar water at higher concentration is more difficult.

By the way, diameters of the antibacterial cycles increase with the concentration of terpinen-4-ol in the essential oil, with the exception of *Staphylococcus aureus* and *Pseudomonas aeruginosa* at very high (98%) concentration of terpinen-4-ol. It may be caused by the less composition of other components in the essential oil and the different cell structures of different types of bacteria.

### **3.2.Quantitative investigation of antibacterial activity by microorganism counting method**

To quantitatively investigate the antibacterial activity of different concentrations of essential oil, microorganism counting method has been applied. Three concentrations (0.5%, 0.25% and 0.1% v/v) have been tested to every chosen type of bacterium. The results are shown in Tables 2,3,4,5 for *Escherichia coli*, *Staphylococcus aureus*, *Candida albicans* and *Pseudomonas aeruginosa*, respectively.

It is clear from experimental results that with 0.5% and 0.25% v/v, all fractions of the essential oil have the antibacterial efficiency of about 90%, including of the fraction of 98% terpinen-4-ol to all chosen types of bacteria, except the little lower efficiency for the case of *Pseudomonas aeruginosa*. This means that the method is more quantitative than the hole boring method, where the non-polar essential oil is less diffusive in polar water media.

With the concentration of 0.1% v/v, the antibacterial effect is still rather high to *Escherichia coli* (more than 80%), *Staphylococcus aureus* (more than 85%) and *Candida albican* (more than 90%). The high antibacterial activity of the fractions of lower terpinen-4-ol composition points out that beside terpinen-4-ol, other components in the tea tree oil have also the added effect of antibacterium.

### **3.3.Investigation of tea tree oil application in cosmetics**


Since tea tree oil is not dissolved in water, the emulsifying reagent Tween-80 is applied with the concentration of 1%. Two concentrations of essential oil have been used: 0.25% and 0.1%. The tests have been carried out to body soap and shampoo. Investigation results are shown in Tables 6,7,8,9.

It has been shown that with only 0.1% of the essential oil, the product is kept with pleasant odor and no colour-change.

## **4.CONCLUSION**

Through the experimental results, it could be pointed out:

- Tea tree oil has strong antibacterial activity to *Escherichia coli*, *Staphylococcus aureus* and *Candida albicans*, and a little weaker to *Pseudomonas aeruginosa*.
- The more concentration of terpinen-4-ol in the essential oil is, the more effective its antibacterial activity is. Nevertheless, not only terpinen-4-ol but also other components in the tea tree oil have the antibacterial effect.
- The accepted dose of all fractions of the tea tree oil in practical use is 0.25%.
- By adding 1% of the emulsifying reagent Tween-80, 0.25% of the tea tree oil could be applied in the cosmetic production for antibacterial purpose.



**Figure 1.** Dependence of the antibacterial cycle diameter on the used amount of essential oil  
 a) *Escherichia coli*, b) *Staphylococcus aureus*, c) *Candida albicans*, d) *Pseudomonas aeruginosa*

**Table 1.** Antibacterial effect of essential oil diffusion and evaporation

Used amount of essential oil (μl)	Diffusion only	Evaporation only	Both diffusion and evaporation
	Diameter of the antibacterial cycle (mm)		
25	19	7	19
25	19	7	19
25	19	7	19
Average			
25	19	7	19

**Table 2.** Investigation result for antibacterial activity to *Escherichia coli*

Sample	Concentration (v/v)	Bacterium before contact (cfu/ml)	Alive bacterium (cfu/ml)	Percentage of killed bacteria (%)
Original essential oil	0.1	$5.10^6$	882.000	83.3
	0.25	$5.10^6$	640.000	87.2
	0.5	$5.10^6$	375.000	92.5
Fraction of 25% Terpinen-4-ol	0.1	$5.10^6$	744.000	80.4
	0.25	$5.10^6$	800.000	78.9
	0.5	$5.10^6$	546.000	85.6
Fraction of 50% Terpinen-4-ol	0.1	$5.10^6$	594.000	84.4
	0.25	$5.10^6$	480.000	87.4
	0.5	$5.10^6$	276.000	92.7

Fraction of 75% Terpinen-4-ol	0.1	$3.8.10^6$	516.000	86.4
	0.25	$3.8.10^6$	468.000	87.7
	0.5	$3.8.10^6$	246.000	93.5
Fraction of 90% Terpinen-4-ol	0.1	$3.8.10^6$	450.000	88.2
	0.25	$3.8.10^6$	392.000	89.7
	0.5	$5.6.10^6$	351.000	93.7
Fraction of 98% Terpinen-4-ol	0.1	$5.6.10^6$	648.000	88.4
	0.25	$5.6.10^6$	396.000	92.9
	0.5	$5.6.10^6$	333.000	94.1

**Table 3.** Investigation result for antibacterial activity to *Staphylococcus aureus*

Sample	Concentration (v/v)	Bacterium before contact (cfu/ml)	Alive bacterium (cfu/ml)	Percentage of killed bacteria (%)
Original essential oil	0.1	$3.02.10^7$	2.160.000	92.8
	0.25	$3.02.10^7$	1.383.000	95.4
	0.5	$3.02.10^7$	826.000	97.3
Fraction of 25% Terpinen-4-ol	0.1	$1.62.10^7$	2.032.000	87.5
	0.25	$1.62.10^7$	549.000	96.6
	0.5	$1.62.10^7$	350.000	97.8
Fraction of 50% Terpinen-4-ol	0.1	$1.62.10^7$	231.600	85.7
	0.25	$1.62.10^7$	504.000	96.9
	0.5	$1.62.10^7$	312.000	98.1
Fraction of 75% Terpinen-4-ol	0.1	$8.8.10^6$	795.000	91.0
	0.25	$8.8.10^6$	729.000	91.7
	0.5	$8.8.10^6$	454.000	94.8
Fraction of 90% Terpinen-4-ol	0.1	$8.8.10^6$	1.450.000	83.5
	0.25	$8.8.10^6$	795.000	91.0
	0.5	$8.8.10^6$	382.000	95.7
Fraction of 98% Terpinen-4-ol	0.1	$3.02.10^7$	2.630.000	91.3
	0.25	$3.02.10^7$	2.240.000	92.6
	0.5	$3.02.10^7$	1.130.000	96.3

**Table 4.** Investigation result for antibacterial activity to *Pseudomonas aeruginosa*

Sample	Concentration (v/v)	Bacterium before contact (cfu/ml)	Alive bacterium (cfu/ml)	Percentage of killed bacteria (%)
Original essential oil	0.1	$2.32.10^7$	7.080.000	69.5
	0.25	$2.32.10^7$	3.300.000	85.8
	0.5	$2.32.10^7$	3.024.000	87.0

Fraction of 25% Terpinen-4-ol	0.1	$2.32.10^7$	4.920.000	78.8
	0.25	$2.32.10^7$	2.120.000	90.9
	0.5	$2.32.10^7$	1.085.000	95.3
Fraction of 50% Terpinen-4-ol	0.1	$2.32.10^7$	5.784.000	75.1
	0.25	$2.32.10^7$	3.140.000	86.5
	0.5	$2.32.10^7$	1.764.000	92.4
Fraction of 75% Terpinen-4-ol	0.1	$2.32.10^7$	5.760.000	75.7
	0.25	$2.32.10^7$	3.220.000	86.1
	0.5	$2.32.10^7$	1.764.000	92.4
Fraction of 90% Terpinen-4-ol	0.1	$1.12.10^7$	3.696.000	67.0
	0.25	$1.12.10^7$	3.159.000	71.79
	0.5	$1.12.10^7$	1.750.000	84.4
Fraction of 98% Terpinen-4-ol	0.1	$1.12.10^7$	4.074.000	63.6
	0.25	$1.12.10^7$	3.636.000	67.5
	0.5	$1.12.10^7$	2.580.000	77.0

**Table 5.** Investigation result for antibacterial activity to *Candida albicans*

Sample	Concentration (v/v)	Bacterium before contact (cfu/ml)	Alive bacterium (cfu/ml)	Percentage of killed bacteria (%)
Original essential oil	0.1	$7.2.10^6$	588.000	91.9
	0.25	$7.2.10^6$	500.000	93.1
	0.5	$7.2.10^6$	320.000	95.6
Fraction of 25% Terpinen-4-ol	0.1	$7.2.10^6$	720.000	90.0
	0.25	$7.2.10^6$	580.000	91.9
	0.5	$7.2.10^6$	224.000	96.9
Fraction of 50% Terpinen-4-ol	0.1	$7.2.10^6$	552.000	92.3
	0.25	$7.2.10^6$	280.000	96.2
	0.5	$7.2.10^6$	200.000	97.2
Fraction of 75% Terpinen-4-ol	0.1	$5.8.10^6$	444.000	92.4
	0.25	$5.8.10^6$	330.000	94.3
	0.5	$5.8.10^6$	280.000	95.2
Fraction of 90% Terpinen-4-ol	0.1	$5.8.10^6$	396.000	93.2
	0.25	$5.8.10^6$	300.000	94.9
	0.5	$5.8.10^6$	240.000	95.9
Fraction of 98% Terpinen-4-ol	0.1	$1.4.10^7$	456.000	96.7
	0.25	$1.4.10^7$	450.000	96.8
	0.5	$1.4.10^7$	288.000	98.9

**Table 6.** Investigation result for antibacterial activity of the essential oil in cosmetics to *E.coli*

Sample	Concentration (v/v)	Bacterium before contact (cfu/ml)	Alive bacterium (cfu/ml)	Percentage of killed bacteria (%)
Body soap	0	$3.12.10^7$	5.000.000	84.0
	0.1	$3.12.10^7$	1.048.000	96.6

	0.25	$3.12.10^7$	4.560.00	98.5
Shampoo	0	$3.12.10^7$	5.920.000	81.0
	0.1	$3.12.10^7$	1.144.000	96.3
	0.25	$3.12.10^7$	520.000	98.3

**Table 7.** Investigation result for antibacterial activity of the essential oil in cosmetics to *Staphylococcus aureus*

Sample	Concentration (v/v)	Bacterium before contact (cfu/ml)	Alive bacterium (cfu/ml)	Percentage of killed bacteria (%)
Body soap	0	$2.24.10^7$	5.000.000	77.78
	0.1	$2.24.10^7$	93.000	95.8
	0.25	$2.24.10^7$	496.000	97.8
Shampoo	0	$2.24.10^7$	5.920.000	73.6
	0.1	$2.24.10^7$	456.000	98.0
	0.25	$2.24.10^7$	336.000	98.5

**Table 8.** Investigation result for antibacterial activity of the essential oil in cosmetics to *Candida albicans*

Sample	Concentration (v/v)	Bacterium before contact (cfu/ml)	Alive bacterium (cfu/ml)	Percentage of killed bacteria (%)
Body soap	0	$6.6.10^6$	1.480.000	77.6
	0.1	$6.6.10^6$	420.000	93.6
	0.25	$6.6.10^6$	330.000	95.0
Shampoo	0	$6.6.10^6$	1.560.000	76.4
	0.1	$6.6.10^6$	480.000	92.7
	0.25	$6.6.10^6$	344.000	94.8

**Table 9.** Investigation result for antibacterial activity of the essential oil in cosmetics to *Pseudomonas aeruginosa*

Sample	Concentration (v/v)	Bacterium before contact (cfu/ml)	Alive bacterium (cfu/ml)	Percentage of killed bacteria (%)
Body soap	0	$2.8.10^7$	6.960.000	75.7
	0.1	$2.8.10^7$	2.808.000	90.2

	0.25	$2.8.10^7$	1.152.000	96.0
Shampoo	0	$2.8.10^7$	7.080.000	75.2
	0.1	$2.8.10^7$	2.832.000	90.1
	0.25	$2.8.10^7$	1.248.000	95.6

## NGHIÊN CỨU ĐẶC TÍNH KHÁNG KHUẨN CỦA TINH DẦU TRÀM TRÀ VÀ KHẢ NĂNG ỨNG DỤNG TRONG MỸ PHẨM

**Phan Đình Tuấn, Hoàng Minh Nam, Nguyễn Thị Thanh Nga**  
Trường Đại Học Bách Khoa, ĐHQG-HCM

**TÓM TẮT:** Bài báo trình bày các kết quả nghiên cứu về đặc tính kháng khuẩn của tinh dầu trà m trà giống Úc trồng tại Đồng Tháp Mười của Đồng Bằng Sông Cửu Long và khả năng sử dụng nó trong sản xuất các mỹ phẩm. Kết quả nghiên cứu chỉ ra rằng không chỉ có terpinen 4-ol mà các cấu tử khác có mặt trong tinh dầu trà m trà đều có khả năng kháng khuẩn. Với hàm lượng sử dụng 0,25%, các phân đoạn khác nhau của tinh dầu trà m trà đều có khả năng tiêu diệt các chủng *Escherichia coli*, *Staphylococcus aureus*, *Pseudomonas aeruginosa* và *Candida albicans* là các chủng điển hình đến 90%. Việc ứng dụng thử nghiệm các phân đoạn khác nhau của tinh dầu trà m trà trong sản xuất mỹ phẩm cho thấy với hàm lượng 0,25%, hoạt tính kháng khuẩn của tinh dầu luôn được đảm bảo mà vẫn có mùi dễ chịu. Kết quả này góp phần củng cố triển vọng ứng dụng của tinh dầu trà m trà trong việc sản xuất các chất tẩy rửa, mỹ phẩm và dược phẩm.

### REFERENCES

- [1]. Lê Ngọc Thạch, *Tinh dầu*, NXB Đại Học Quốc Gia Tp Hồ Chí Minh, (2003)
- [2]. Vương Ngọc Chính, *Hương Liệu Mỹ Phẩm*, NXB Đại Học Quốc Gia Tp Hồ Chí Minh, (2005)
- [3]. Nguyễn Đức Lượng, Phan Thị Huyền, Nguyễn Ánh Tuyết, *Thí Nghiệm Vi Sinh Vật học*, NXB Đại Học Quốc Gia Tp Hồ Chí Minh, (2006).
- [4]. Đỗ Tất Lợi, *Những cây thuốc và Vị thuốc Việt Nam*, NXB Khoa Học và Kỹ Thuật, Hà Nội, (1986).
- [5]. Đỗ Tất Lợi, *Tinh Dầu Việt Nam*, NXB Y Học, (1985).
- [6]. Vũ Ngọc Lộ và nhiều tác giả, *Những cây tinh dầu Việt Nam: Khai Thác, chế biến, Ứng Dụng*, NXB Khoa học và Kỹ thuật, Hà Nội, (1996).
- [7]. Nguyễn Lâm Dũng, *Một số Phương Pháp nghiên cứu vi sinh vật học*, Tập II, NXB Khoa học và Kỹ Thuật, Hà Nội, (1976).
- [8]. Louis Hồ Tấn Tài, *Các sản phẩm tẩy rửa và chăm sóc cá nhân- Lý thuyết và Ứng Dụng*. Công Ty Liksin, TP HCM, (1999).
- [9]. Lê Nhất Thông, *Nghiên cứu công nghệ tách thu hồi tinh dầu Trà m Trà giống Úc (*Melaleuca Alternifolia*)trồng trên đất phèn Đồng Bằng Sông Cửu Long, Việt Nam*, Luận văn Thạc Sĩ, trường Đại Học Bách Khoa Tp.HCM, (1999)
- [10]. *The Essential oil*, Vol 1,2,3,4, New York, (1982).
- [11]. J E Drinnan B.Agar, *Development of the North Queensland Tea Tree Industry*, (1998)..

- [12]. C F Carson, T V Riley, *Antimicrobial Activity of Tea Tree Oil*, DPIE Copy Centre, (1998).
- [13]. Southwell et al., *Quality assurance for Tea Tree Oil safety investigative samples* , Canprint, (2006).
- [14]. C F Carson, T V Riley, *Antimicrobial anti-inflammatory of Tea Tree Oil*, DPIE Copy Centre, (2000).
- [15]. Jane Southwell et al, *Metabolism of 1,8- cineol in Tea Tree (Melaleuca Alternifolia) by pygo Beetle*, (1994).
- [16]. Diane M. Martin, Joorg Bohlmann, *Phytochemistry* 65, (2004), p.1223–1229.
- [17]. S.D. Cox et al., *The mode of antimicrobial action of the essential oil of Melaleuca alternifolia (tea tree oil)*, *Journal of Applied Microbiology*, (2000), 88, p.170–175.
- [18]. K.A. Hammer, C.F. Carson and T.V. Riley, *Antifungal activity of the components of Melaleuca alternifolia (tea tree) oil*, *Journal of Applied Microbiology*, (2003), 95, p.853–860.
- [19]. Shelley F. Walton, *Acaricidal Activity of Melaleuca alternifolia (Tea Tree) Oil*, *Arch Dermatol/ Vol. 140, May 2004*, p.563-567.
- [20]. S.D. Cox, C.M. Mann and J.L. Markham, *Interactions between components of the essential oil of Melaleuca alternifolia*, *Journal of Applied Microbiology* (2001), 91,p.492-497.
- [21]. Christine F. Carson, Brian J. Mee, and Thomas V. Riley, *Mechanism of Action of Melaleuca alternifolia (Tea Tree) Oil on Staphylococcus aureus, determined by Time-Kill, Lysis, Leakage, and Salt Tolerance Assays and Electron Microscopy*, *Antimicrobial agents and Chemotherapy*, June 2002, p.1914–1920.
- [22]. S.A. Burt and R.D. Reinders, *Antibacterial activity of selected plant essential oils against Escherichia coli*, *Letters in Applied Microbiology*,(2003), 36, p.162–167.
- [23]. J.E Gustafson et al., *Effect of tee tree oil on Escherichia coli*, *Letters in Applied Microbiology*, (1998), 26, p.194-198.