

Research Article

STRONG CONVERGENCE OF A HYBRID ITERATION FOR GENERALIZED MIXED EQUILIBRIUM PROBLEM AND BREGMAN TOTALLY QUASI-ASYMPTOTICALLY NONEXPANSIVE MAPPING IN BANACH SPACES

Nguyen Trung Hieu

Department of Mathematics and Information Technology Teacher Education,
Dong Thap University, Cao Lanh City, Dong Thap Province, Viet Nam

*Corresponding author: Nguyen Trung Hieu, Email: ngtrunghieu@dtu.edu.vn

Received: June 20, 2021; Revised: August 20, 2021; Accepted: September, 2021

ABSTRACT

The purpose of this paper is to combine the Bregman distance with the shrinking projection method to introduce a new hybrid iteration process for a generalized mixed equilibrium problem and a Bregman totally quasi-asymptotically nonexpansive mapping. After that, under some suitable conditions, we prove that the proposed iteration strongly converges to the Bregman projection of the initial point onto common element set of the solution set of a generalized mixed equilibrium problem and the fixed point set of a Bregman totally quasi-asymptotically nonexpansive mapping in reflexive Banach spaces. This theorem extends and improves the results in (Alizadeh & Moradlou, 2016) from a generalized hybrid mapping and an equilibrium problem in Hilbert spaces to a Bregman totally quasi-asymptotically nonexpansive mapping and a generalized mixed equilibrium problem in reflexive Banach spaces. The obtained result is applied to a generalized mixed equilibrium problem and a Bregman quasi-asymptotically nonexpansive mapping in reflexive Banach spaces. In addition, an example is provided to illustrate for the proposed iteration process.

Keywords: Bregman totally quasi-asymptotically nonexpansive mapping; generalized mixed equilibrium problem; hybrid iteration process; reflexive Banach spaces

1. Introduction and preliminaries

Suppose that X is a real reflexive Banach space, Ω is a nonempty, closed and convex subset of X , X^* is a the dual space of X . Let $f : \Omega \times \Omega \rightarrow \mathbb{R}$, $\varphi : \Omega \rightarrow \mathbb{R}$ be two function and $\psi : \Omega \rightarrow X^*$ be a mapping. We denote the value of $u^* \in X^*$ at $u \in X$ by $\langle u^*, u \rangle$. The generalized mixed equilibrium problem (GMEP) is to find $u \in \Omega$ such that $f(u, v) + \langle \psi(u), v - u \rangle + \varphi(v) \geq \varphi(u)$ for all $v \in \Omega$. The set of solutions of (GMEP) is denoted by $GMEP(f, \varphi, \psi) = \{u \in \Omega : f(u, v) + \langle \psi(u), v - u \rangle + \varphi(v) \geq \varphi(u), \forall v \in \Omega\}$. Note that, if $\varphi \equiv 0$ and $\psi \equiv 0$, the problem (GMEP) is reduced into the equilibrium problem (EP) which is to find $u \in \Omega$ such that $f(u, v) \geq 0$ for all $v \in \Omega$.

Cite this article as: Nguyen Trung Hieu (2021). Using creative methodology to explore factors influencing teacher educator identity. *Ho Chi Minh City University of Education Journal of Science*, 18(9), 1359-1367.

In recent times, there were many methods for solving the above problems. In 2016, Darvish introduced an iterative method for finding common elements of the solutions set of the problem (GMEP) and the fixed points set of a Bregman strongly nonexpansive mapping in reflexive Banach spaces. In 2016, Zhu and Huang introduced a new hybrid iterative scheme for finding common solutions of the problem (EP) and fixed points of Bregman totally quasi-asymptotically nonexpansive mappings. In 2018, Ni and Wen proposed a new iterative scheme for finding a common solution of a system of the problem (GMEP) and fixed points of a finite family of Bregman totally quasi-asymptotically nonexpansive mappings. Note that these convergence results extend and improve the existing results from Hilbert spaces or smooth Banach spaces to reflexive Banach spaces. Therefore, an interesting work naturally raised is to continue to generalize the existing convergence results from Hilbert spaces to reflexive Banach spaces.

In this paper, motivated by the iteration process in (Alizadeh & Moradlou, 2016), we introduce a new hybrid iterative scheme which is to find common elements of the set of solutions of the problem (GMEP) and the set of fixed points of Bregman totally quasi-asymptotically nonexpansive mappings. After that, we prove a strong convergence theorem for the proposed iteration in reflexive Banach spaces. In addition, we give a numerical example to illustrate the obtained results.

Now, we recall some notions and results which will be useful in what follows.

Assume that $g : X \rightarrow (-\infty, +\infty]$ is a lower semi-continuous, convex and proper function. We denote the domain of g by $\text{dom } g = \{u \in X : g(u) < +\infty\}$. For any $u \in \text{int}(\text{dom } g)$ and $v \in X$, we denote by $g'(u, v) = \lim_{\lambda \rightarrow 0^+} \frac{g(u + \lambda v) - g(u)}{\lambda}$ (1.1) the right-hand derivative of g at u in the direction v . The function g is called *Gâteaux differentiable at u* if the limit (1.1) exists for all v . Then the gradient of g at u is $\nabla g(u)$, which is defined by $\langle \nabla g(u), v \rangle = g'(u, v)$ for all $v \in X$. The function g is called *Fréchet differentiable at u* if the limit (1.1) is attained uniformly in $\|v\| = 1$. The function g is called *uniformly Fréchet differentiable on a subset Ω of X* if the limit (1.1) is attained uniformly for $u \in \Omega$ and $\|v\| = 1$.

Note that if g is uniformly Fréchet differentiable, then g is uniformly continuous (see [Ambrosetti & Prodi, 1993, Theorem 1.8]). If g is Gâteaux differentiable and lower semi-continuous convex, then g is bounded on bounded sets if and only if ∇g is bounded on bounded sets (see [Ambrosetti & Prodi, 1993, Proposition 1.1.11]). Furthermore, if g is uniformly Fréchet differentiable and bounded on bounded subsets, then ∇g is uniformly continuous on bounded subsets of X^* (see [Reich & Sabach, 2009, Proposition 1]).

Let $u \in \text{int}(\text{dom } g)$, the Fenchel conjugate of g is the function $g^* : X^* \rightarrow (-\infty, +\infty]$ defined by $g^*(u^*) = \sup\{\langle u^*, u \rangle - g(u) : u \in X\}$ for all $u^* \in X^*$.

Definition 1.1 [Chang *et al.*, 2014, Definition 2.2]. Let X be a real reflexive Banach space and $g : X \rightarrow (-\infty, +\infty]$ be a function. Then g is called *Legendre* if

(L1) $\text{int}(\text{dom} g) \neq \emptyset$, g is Gâteaux differentiable on $\text{int}(\text{dom} g)$ and

$\text{dom}(\nabla g) = \text{int}(\text{dom} g)$.

(L2) $\text{int}(\text{dom} g^*) \neq \emptyset$, g^* is Gâteaux differentiable on $\text{int}(\text{dom} g^*)$ and

$\text{dom}(\nabla g^*) = \text{int}(\text{dom} g^*)$.

Remark 1.2. [Chang *et al.*, 2014, Remark 2.3]. Let X be a real reflexive Banach space and $g : E \rightarrow (-\infty, +\infty]$ be Legendre. Then

(1) g is Legendre if and only if g^* is Legendre.

(2) $\nabla g = (\nabla g^*)^{-1}$, $\text{ran}(\nabla g) = \text{dom}(\nabla g^*)$ and $\text{ran}(\nabla g^*) = \text{dom}(\nabla g) = \text{int}(\text{dom} g)$, where $\text{ran}(\nabla g)$ is the range of ∇g .

Definition 1.3. [Censor & Lent, 1981, p.324]. Let X be a real reflexive Banach space and $g : X \rightarrow (-\infty, +\infty]$ be Gâteaux differentiable. Then $D_g : \text{dom} g \times \text{int}(\text{dom} g) \rightarrow [0, \infty)$, defined by $D_g(u, v) = g(u) - g(v) - \langle \nabla g(v), u - v \rangle$ is called the *Bregman distance* with respect to g .

From the definition, we have $D_g(u, v) + D_g(v, w) - D_g(u, w) = \langle \nabla g(w) - \nabla g(v), u - v \rangle$ for all $u \in \text{dom} g$ and $v, w \in \text{int}(\text{dom} g)$.

Let $g : X \rightarrow (-\infty, +\infty]$ be Gâteaux differentiable and $V_g : X \times X^* \rightarrow [0, \infty)$ be defined by $V_g(u, u^*) = g(u) - \langle u^*, u \rangle + g^*(u^*)$ for all $u \in X$ and $u^* \in X^*$.

Remark 1.4. Let $g : X \rightarrow (-\infty, +\infty]$ be a Gâteaux differentiable function. Then

(1) [Kohsaka & Takahashi, 2005, Lemma 3.2] For any $u \in X$ and $u^* \in X^*$, we have

$$V_g(u, u^*) = D_g(u, \nabla g^*(u^*)).$$

(2) [Kumam *et al.*, 2016, p.7] V_g is convex in the second variable. Furthermore, for

any $u \in \text{dom} g$, $\{u_k\}_{k=1}^m \subset \text{int}(\text{dom} g)$ and $\{t_k\}_{k=1}^m \subset [0, 1]$ with $\sum_{k=1}^m t_k = 1$, we have

$$D_g(u, \nabla g^*(\sum_{k=1}^m t_k \nabla g(u_k))) \leq \sum_{k=1}^m t_k D_g(u, u_k).$$

Definition 1.5. [Butnariu & Iusem, 2000, p.69]. Let X be a real reflexive Banach space, $g : X \rightarrow (-\infty, +\infty]$ be Legendre and Ω be a nonempty, convex and closed subset of $\text{int}(\text{dom} g)$. The *Bregman projection* of $u \in \text{int}(\text{dom} g)$ onto Ω is the unique vector $P_\Omega^g(u) \in \Omega$ satisfying $D_g(P_\Omega^g(u), u) = \inf\{D_g(v, u) : v \in \Omega\}$.

Definition 1.6. [Resmerita, 2004, p.1]. Let X be a real reflexive Banach space and $g : X \rightarrow (-\infty, +\infty]$ be Gâteaux differentiable. Then

(1) g is called *totally convex* at $u \in \text{int}(\text{dom} g)$ if any $\varepsilon > 0$, we have

$$v_g(u, \varepsilon) = \inf\{D_g(v, u) : v \in \text{dom} g, \|v - u\| = \varepsilon\} > 0.$$

(2) g is called *totally convex* if g is totally convex at every point $u \in \text{int}(\text{dom}f)$.

(3) g is called *totally convex on bounded subsets of X* if any nonempty bounded subset E of X and $t > 0$, we have $v_g(E, \varepsilon) = \inf\{v_g(u, \varepsilon) : u \in E \cap \text{dom} g\} > 0$.

Proposition 1.7. [Resmerita, 2004, Proposition 2.2]. *Let X be a real reflexive Banach space, and $g : X \rightarrow \mathbb{R}$ be Gâteaux differentiable. Then g is totally convex at $u \in X$ if and only if for any sequence $\{v_n\} \subset X$ such that $\lim_{n \rightarrow \infty} D_g(v_n, u) = 0$, we have $\lim_{n \rightarrow \infty} \|v_n - u\| = 0$.*

Proposition 1.8. [Butnariu & Iusem, 2000, Lemma 2.1.2]. *Let X be a real reflexive Banach space, and $g : X \rightarrow \mathbb{R}$ be convex and Gâteaux differentiable. Then g is totally convex on bounded sets if and only if for any sequence $\{u_n\}, \{v_n\} \subset X$ such that $\{u_n\}$ is bounded and $\lim_{n \rightarrow \infty} D_g(v_n, u_n) = 0$, we have $\lim_{n \rightarrow \infty} \|v_n - u_n\| = 0$.*

Proposition 1.9. [Butnariu & Resmerita, 2006, Corollary 4.4]. *Let X be a real reflexive Banach space, $g : X \rightarrow (-\infty, +\infty]$ be a Gâteaux differentiable function and totally convex on $\text{int}(\text{dom}g)$, Ω be a nonempty, closed and convex subset and $u \in \text{int}(\text{dom}g)$. Then*

(1) $w = P_\Omega^g(u)$ if and only if $\langle \nabla g(u) - \nabla g(w), w - v \rangle \geq 0$ for all $v \in \Omega$.

(2) $D_g(v, P_\Omega^g(u)) + D_g(P_\Omega^g(u), u) \leq D_g(v, u)$ for all $v \in \Omega$.

Proposition 1.10. *Let X be a real reflexive Banach space and $g : X \rightarrow \mathbb{R}$ be a function.*

(1) [Reich & Sabach, 2010, Lemma 1]. *If g is Gâteaux differentiable and totally convex on X , $u \in X$ and $\{u_n\} \subset X$ satisfying $\{D_g(u_n, u)\}$ is bounded, then the sequence $\{u_n\}$ is bounded.*

(2) [Sabach, 2011, Proposition 2.3]. *If g is Legendre such that ∇g^* is bounded on bounded subsets, $u \in X$ and $\{u_n\} \subset X$ satisfying $\{D_g(u, u_n)\}$ is bounded, then the sequence $\{u_n\}$ is bounded.*

Definition 1.11. [Zalinescu, 2002, p.203, p.207, p.221]. *Let X be a Banach space. We denote by $S_1 = \{u \in X : \|u\| < 1\}$ and $B_\varepsilon = \{u \in X : \|u\| \leq \varepsilon\}$ for some $\varepsilon > 0$. Then*

(1) $g : X \rightarrow \mathbb{R}$ is called *uniformly convex on bounded subsets* if $\rho_\varepsilon(\lambda) > 0$ for all $\lambda, \varepsilon > 0$, where the function $\rho_\varepsilon : [0, \infty) \rightarrow [0, \infty)$ is defined by

$$\rho_\varepsilon(\lambda) = \inf_{u, v \in B_\varepsilon, \|u-v\|=\lambda, \delta \in (0,1)} \frac{\delta g(u) + (1-\delta)g(v) - g(\delta u + (1-\delta)v)}{\delta(1-\delta)}.$$

(2) $g : X \rightarrow \mathbb{R}$ is called *uniformly smooth on bounded subsets* if $\lim_{\lambda \rightarrow 0} \frac{\sigma_\varepsilon(\lambda)}{\lambda} = 0$ for all $\varepsilon > 0$, where the function $\sigma_\varepsilon : [0, \infty) \rightarrow [0, \infty)$ is defined by

$$\sigma_\varepsilon(\lambda) = \sup_{u \in B_\varepsilon, v \in S_1, \delta \in (0,1)} \frac{\delta g(u + (1-\delta)\lambda v) + (1-\delta)g(u - \delta\lambda v) - g(u)}{\delta(1-\delta)}.$$

Remark 1.12. [Naraghirad & Yao, 2013, p.7]. *The function g is uniformly convex on bounded subsets if and only if g is totally convex on bounded subsets.*

Definition 1.13. [Kohsaka and Takahashi, 2005, p.509]. Let X be a Banach space. Then

$g : X \rightarrow (-\infty, +\infty]$ is called *strongly coercive* if $\lim_{\|u\| \rightarrow +\infty} \frac{g(u)}{\|u\|} = +\infty$.

Proposition 1.14. [Zalinescu, 2002, Proposition 3.6.3]. *Let X be a real reflexive Banach space, $g : X \rightarrow \mathbb{R}$ be strongly coercive, continuous and convex. Then g is bounded on bounded subsets and uniformly smooth on bounded subsets if and only if $\text{dom}(g^*) = X^*$, g^* is strongly coercive and uniformly convex on bounded subsets.*

Proposition 1.15. [Zalinescu, 2002, Proposition 3.6.4]. *Let X be a real reflexive Banach space, $g : X \rightarrow \mathbb{R}$ be convex, continuous and bounded on bounded subsets of X . Then the following statements are equivalent.*

- (1) g is uniformly convex on bounded subsets and strongly coercive.
- (2) $\text{Dom}(g^*) = X^*$, g^* is bounded and uniformly smooth on bounded subsets.
- (3) $\text{Dom}(g^*) = X^*$, g^* is Fréchet differentiable and ∇g^* is uniformly continuous on bounded subsets.

Lemma 1.16. [Naraghirad & Yao, 2013, Lemma 2.2]. *Let X be a Banach space, $r > 0$ and $g : X \rightarrow \mathbb{R}$ be convex and uniformly convex on bounded subsets. Then*

$$g\left(\sum_{n=1}^m a_n u_n\right) \leq \sum_{n=1}^m a_n g(u_n) - a_i a_j \rho_\varepsilon(\|u_i - u_j\|)$$

with $i, j \in \{1, 2, \dots, m\}$, $u_n \in B_\varepsilon = \{u \in X : \|u\| \leq \varepsilon\}$ and $a_n \in [0, 1]$ such that $\sum_{n=1}^m a_n = 1$,

and the function ρ_ε is defined as in Definition 1.11.

We denote by $F(S) = \{w \in \Omega : Sw = w\}$ the set of fixed points of $S : \Omega \rightarrow \Omega$.

Definition 1.17. [Chang et al., 2014, Definition 2.10]. Let X be a reflexive Banach space, Ω be a nonempty subset of X , $S : \Omega \rightarrow \Omega$ be a mapping and D_g be the Bregman distance.

Then

- (1) S is called a *Bregman quasi-asymptotically nonexpansive mapping* if $F(S) \neq \emptyset$ and there exists a real sequence $\{\delta_n\} \subset [1, \infty)$ with $\lim_{n \rightarrow \infty} \delta_n = 1$ such that

$$D_g(u, S^n v) \leq \delta_n D_g(u, v) \text{ for all } v \in \Omega \text{ and } u \in F(S).$$

- (2) S is called a *Bregman totally quasi-asymptotically nonexpansive mapping* if $F(S) \neq \emptyset$ and there exist nonnegative real sequences $\{\alpha_n\}, \{\beta_n\}$ with $\lim_{n \rightarrow \infty} \alpha_n = \lim_{n \rightarrow \infty} \beta_n = 0$ and a strictly increasing continuous function $\zeta : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ with $\zeta(0) = 0$ such that

$$D_g(u, S^n v) \leq D_g(u, v) + \alpha_n \zeta(D_g(u, v)) + \beta_n \text{ for all } v \in \Omega \text{ and } u \in F(S).$$

(3) S is called a *Bregman firmly nonexpansive mapping* if

$$\langle \nabla g(Su) - \nabla g(Sv), Su - Sv \rangle \leq \langle \nabla g(u) - \nabla g(v), Su - Sv \rangle \text{ for all } u, v \in \Omega.$$

(4) S is called a *Bregman quasi-nonexpansive mapping* if $F(S) \neq \emptyset$ and

$$D_g(u, Sv) \leq D_g(u, v) \text{ for all } v \in \Omega \text{ and } u \in F(S).$$

Remark 1.18. [Chang et al., 2014, p.42].

(1) If S is a Bregman quasi-asymptotically nonexpansive mapping, then S is a Bregman totally quasi-asymptotically nonexpansive mapping with $\zeta(\lambda) = \lambda$ for all $\lambda \geq 0$, $\alpha_n = \delta_n - 1$ with $\delta_n \geq 1$ satisfying $\lim_{n \rightarrow \infty} \delta_n = 1$ and $\beta_n = 0$; but the converse is not true.

(2) If S is a Bregman firmly nonexpansive mapping, then S is a Bregman quasi-nonexpansive mapping.

Definition 1.19. [Zhu & Huang, 2016, Definition 2.10]. Let X be a Banach space, Ω be a nonempty subset of X , $S : \Omega \rightarrow \Omega$ be a mapping. Then

(1) S is called *closed* if any sequence $\{u_n\}$ in Ω such that $\lim_{n \rightarrow \infty} u_n = u \in \Omega$ and $\lim_{n \rightarrow \infty} Su_n = v \in \Omega$, we have $Su = v$.

(2) S is called *uniformly asymptotically regular* on Ω if for all bounded subset U of Ω we have $\lim_{n \rightarrow \infty} \sup_{x \in U} \|S^{n+1}x - S^n x\| = 0$.

Lemma 1.20. [Chang et al., 2014, Lemma 2.16]. Let X be a real reflexive Banach space, Ω be a nonempty, closed and convex subset of X , $g : X \rightarrow (-\infty, +\infty]$ be a Legendre function which is totally convex on bounded subsets of X , $S : \Omega \rightarrow \Omega$ be a closed and Bregman totally quasi-asymptotically nonexpansive mapping. Then $F(S)$ is convex and closed.

In order to solve (GMEP), we suppose that f satisfies the following hypotheses:

(C1) $f(u, u) = 0$ for all $u \in \Omega$.

(C2) $f(u, v) + f(v, u) \leq 0$ for all $u, v \in \Omega$.

(C3) $\limsup_{\lambda \rightarrow 0} f(\lambda w + (1 - \lambda)u, v) \leq f(u, v)$ for all $u, v, w \in \Omega$,

(C4) For each $u \in \Omega$, $v \mapsto f(u, v)$ is convex and lower semi-continuous.

Definition 1.21. [Darvish, 2016, Definition 2.4]. Let X be a real reflexive Banach space, Ω be a nonempty, convex and closed subset of X . Suppose that $f : \Omega \times \Omega \rightarrow \mathbb{R}$ satisfies (C1)-(C4), $\varphi : \Omega \rightarrow \mathbb{R}$ is convex and lower semi-continuous, $\psi : \Omega \rightarrow X^*$ is continuous monotone. The mixed resolvent of f is the mapping $\text{Res}_{f, \varphi, \psi}^g : X \rightarrow 2^\Omega$ which is defined by

$$\begin{aligned} \text{Res}_{f, \varphi, \psi}^g(u) = \{w \in \Omega : f(w, v) + \varphi(v) + \langle \psi(u), v - w \rangle \\ + \langle \nabla f(w) - \nabla f(u), v - w \rangle \geq \varphi(w), \forall v \in \Omega\}. \end{aligned}$$

Note that if $g : X \rightarrow (-\infty, +\infty]$ is strongly coercive and Gâteaux differentiable, then $\text{dom}(\text{Res}_{f, \varphi, \psi}^g) = X$, see [Darvish, 2016, Lemma 2.7]. We find that the formula of the function $\text{Res}_{f, \varphi, \psi}^g$ contains the term $\psi(u)$ for all $u \in X$. Since $\text{dom} \psi = \Omega \subset X$, the value

$\psi(u)$ does not exist for all $u \in X \setminus \Omega$. Motivated by this confusion, we revise the formula of the function $\text{Res}_{f,\varphi,\psi}^g$ by replacing the term $\psi(u), u \in X$ by $\psi(w), w \in \Omega$. This formula has been stated in (Ni & Wen, 2018, Lemma 2.5), where $\text{Res}_{f,\varphi,\psi}^g$ is denoted by T_r^G as follows.

$$\begin{aligned} \text{Res}_{f,\varphi,\psi}^g(u) = \{w \in \Omega : f(w, v) + \varphi(v) + \langle \psi(w), v - w \rangle \\ + \langle \nabla f(w) - \nabla f(u), v - w \rangle \geq \varphi(w), \forall v \in \Omega\}. \end{aligned} \quad (1.2)$$

The following lemma presents some properties of $\text{Res}_{f,\varphi,\psi}^g$ which is defined by (1.2).

Lemma 1.22. [Ni & Wen, 2018, Lemma 2.5]. *Let X be a real reflexive Banach space, Ω be a nonempty, closed and convex subset of X , $g : X \rightarrow \mathbb{R}$ be Legendre and $f : \Omega \times \Omega \rightarrow \mathbb{R}$ be a bifunctional satisfying (C1)-(C4). Then*

- (1) $\text{Res}_{f,\varphi,\psi}^g$ is a single-valued and Bregman firmly nonexpansive mapping.
- (2) $F(\text{Res}_{f,\varphi,\psi}^g) = \text{GMEP}(f, \varphi, \psi)$, $\text{GMEP}(f, \varphi, \psi)$ is convex and closed.
- (3) For all $u \in X$ and $v \in F(\text{Res}_{f,\varphi,\psi}^g)$, we have

$$D_g(v, \text{Res}_{f,\varphi,\psi}^g(u)) + D_g(\text{Res}_{f,\varphi,\psi}^g(u), u) \leq D_g(v, u).$$

2. Main results

The following result shows the strong convergence of a hybrid iteration process for a generalized mixed equilibrium problem and a Bregman totally quasi-asymptotically nonexpansive mapping in reflexive Banach spaces.

Theorem 2.1. *Let X be a real reflexive Banach space, Ω be a nonempty, closed and convex subset of X , $g : X \rightarrow \mathbb{R}$ be Legendre, strongly coercive, bounded, totally convex and Fréchet differentiable on bounded subsets. Suppose that $f : \Omega \times \Omega \rightarrow \mathbb{R}$ satisfies (C1)-(C4), $\varphi : \Omega \rightarrow \mathbb{R}$ is lower semi-continuous and convex, $\psi : \Omega \rightarrow X^*$ is continuous monotone, $S : \Omega \rightarrow \Omega$ is a closed, uniformly asymptotically regular and Bregman totally quasi-asymptotically nonexpansive mapping with $\{\alpha_n\}, \{\beta_n\} \subset [0, \infty)$ satisfying $\lim_{n \rightarrow \infty} \alpha_n = \lim_{n \rightarrow \infty} \beta_n = 0$ and a strictly increasing continuous function $\zeta : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ with $\zeta(0) = 0$ such that $\mathcal{F} = F(S) \cap \text{GMEP}(f, \varphi, \psi)$ is bounded and nonempty. Let $\{z_n\}$ be a sequence generated by: $z_1 \in \Omega, \Omega_1 = \Omega$ and*

$$\begin{cases} u_n = \nabla g^*(a_n \nabla g(z_n) + (1 - a_n) \nabla g(S^n z_n)) \\ v_n \in \Omega : f(v_n, v) + \varphi(v) + \langle \psi(v_n), v - v_n \rangle + \langle \nabla g(v_n) - \nabla g(z_n), v - v_n \rangle \geq \varphi(v_n), \forall v \in \Omega \\ w_n = \nabla g^*(b_n \nabla g(u_n) + (1 - b_n) \nabla g(S^n v_n)) \\ \Omega_{n+1} = \{u \in \Omega_n : D_g(u, w_n) \leq D_g(u, z_n) + \gamma_n\} \\ z_{n+1} = P_{\Omega_{n+1}}^g(z_1), n \in \mathbb{N}^* \end{cases} \quad (2.1)$$

where $\gamma_n = \alpha_n \sup\{\zeta(D_g(u, z_n)) : u \in \mathcal{F}\} + \beta_n$ and $\{a_n\}, \{b_n\} \subset [0, 1]$ such that $\lim_{n \rightarrow \infty} a_n = 1$ and $\liminf_{n \rightarrow \infty} b_n(1 - b_n) > 0$. Then the sequence $\{z_n\}$ strongly converges to $p = P_{\mathcal{F}}^g(z_1)$.

Proof. We divide the proof of this theorem into six steps.

Step 1. We show that $P_{\mathcal{F}}^g(x_1)$ is well-defined. Indeed, it follows from Lemma 1.20 and Lemma 1.22 that $F(S)$ and $GMEP(f, \varphi, \psi)$ are closed and convex. Therefore, by combining this with the assumption, we obtain that $\mathcal{F} = F(S) \cap GMEP(f, \varphi, \psi)$ is a nonempty, closed and convex subset of Ω . This fact ensures that $P_{\mathcal{F}}^g(z_1)$ is well-defined.

Step 2. We show that $P_{\Omega_{n+1}}^g(z_1)$ is well-defined. We first claim by mathematical induction that Ω_n is convex and closed for all $n \in \mathbb{N}^*$. Obviously, for $n = 1$, we have $\Omega_1 = \Omega$ is closed and convex. Now we suppose that Ω_k is convex and closed for some $k \in \mathbb{N}^*$. Then, by the definition of Ω_{n+1} , we have

$$\Omega_{k+1} = \{u \in \Omega_k : \langle \nabla g(z_k), u - z_k \rangle - \langle \nabla g(w_k), u - w_k \rangle \leq \gamma_k - g(z_k) + g(w_k)\}. \quad (2.2)$$

By combining (2.2) with the continuity of $\nabla g(\cdot)$, we get that Ω_{k+1} is convex and closed. Therefore, Ω_n is convex and closed for all $n \in \mathbb{N}^*$. Next, we will claim by mathematical induction that $\mathcal{F} \subset \Omega_n$ for all $n \in \mathbb{N}^*$. Obviously, we have $\mathcal{F} \subset \Omega = \Omega_1$. Now, we suppose that $\mathcal{F} \subset \Omega_k$ for some $k \in \mathbb{N}^*$. We will show that $\mathcal{F} \subset \Omega_{k+1}$. Indeed, for any $u \in \mathcal{F}$, we get $u \in \Omega_k$. By using Remark 1.4.(2), we have

$$\begin{aligned} D_g(u, u_k) &= D_g(u, \nabla g^*(a_k \nabla g(z_k) + (1 - a_k) \nabla g(S^k z_k))) \leq a_k D_g(u, z_k) + (1 - a_k) D_g(u, S^k z_k) \\ &\leq a_k D_g(u, z_k) + (1 - a_k) [D_g(u, z_k) + \alpha_k \zeta(D_g(u, z_k)) + \beta_k] \\ &= D_g(u, z_k) + (1 - a_k) [\alpha_k \zeta(D_g(u, z_k)) + \beta_k] \leq D_g(u, z_k) + \alpha_k \zeta(D_g(u, z_k)) + \beta_k. \end{aligned} \quad (2.3)$$

Furthermore, by the definition of v_n and Definition 1.21, we have $v_n = \text{Res}_{f, \varphi, \psi}^g(z_n)$. It follows from Remark 1.18 and Lemma 1.22 that $\text{Res}_{f, \varphi, \psi}^g$ is a Bregman quasi-nonexpansive mapping. Therefore $D_g(u, v_k) = D_g(u, \text{Res}_{f, \varphi, \psi}^g(z_k)) \leq D_g(u, z_k)$.

Next, by using Remark 1.4.(2), we obtain

$$\begin{aligned} D_g(u, w_k) &= D_g(u, \nabla g^*(b_k \nabla g(u_k) + (1 - b_k) \nabla g(S^k v_k))) \leq b_k D_g(u, u_k) + (1 - b_k) D_g(u, T^k v_k) \\ &\leq b_k D_g(u, u_k) + (1 - b_k) [D_g(u, v_k) + \alpha_k \zeta(D_g(u, v_k)) + \beta_k]. \end{aligned} \quad (2.5)$$

It follows from (2.4) and the strictly increasing property of ζ that $\zeta(D_g(u, v_k)) < \zeta(D_g(u, z_k))$. Then, from (2.4), (2.5) becomes

$$D_g(u, w_k) \leq b_k D_g(u, u_k) + (1 - b_k) [D_g(u, z_k) + \alpha_k \zeta(D_g(u, z_k)) + \beta_k]. \quad (2.6)$$

By substituting (2.3) into (2.6), we have

$$\begin{aligned} D_g(u, w_k) &\leq b_k [D_g(u, z_k) + \alpha_k \zeta(D_g(u, z_k)) + \beta_k] + (1 - b_k) [D_g(u, z_k) + \alpha_k \zeta(D_g(u, z_k)) + \beta_k] \\ &= D_g(u, z_k) + \alpha_k \zeta(D_g(u, z_k)) + \beta_k \leq D_g(u, z_k) + \gamma_k. \end{aligned} \quad (2.7)$$

This implies that $u \in \Omega_{k+1}$ and hence $\mathcal{F} \subset \Omega_{k+1}$. Therefore, we conclude that $\mathcal{F} \subset \Omega_n$ for all $n \in \mathbb{N}^*$. By the assumption $\mathcal{F} \neq \emptyset$, we obtain $\Omega_{n+1} \neq \emptyset$. Therefore, we find that $P_{\Omega_{n+1}}^g(z_1)$ is well-defined.

Step 3. We show that $\{D_g(z_n, z_1)\}, \{z_n\}$ are bounded and $\lim_{n \rightarrow \infty} D_g(z_n, z_1)$ exists. Indeed, since $z_n = P_{\Omega_n}^g(z_1)$, by Proposition 1.9, we get $D_g(y, z_n) + D_g(z_n, z_1) \leq D_g(y, z_1), \forall y \in \Omega_n$. (2.8)

Let $u \in \mathcal{F}$. Since $\mathcal{F} \subset \Omega_n$, we get $u \in \Omega_n$. By choosing $y = u$ in (2.8), we obtain

$$D_g(u, z_n) + D_g(z_n, z_1) \leq D_g(u, z_1). \quad (2.9)$$

This implies that $D_g(z_n, z_1) \leq D_g(u, z_1) - D_g(u, z_n) \leq D_g(u, z_1)$. Therefore, $\{D_g(z_n, z_1)\}$ is bounded. Then, by Proposition 1.10(1), we conclude that the sequence $\{z_n\}$ is bounded. Furthermore, we have $z_{n+1} = P_{\Omega_{n+1}}^g(z_1) \in \Omega_{n+1} \subset \Omega_n$. By choosing $y = z_{n+1}$ in (2.8), we get $D_g(z_{n+1}, z_n) + D_g(z_n, z_1) \leq D_g(z_{n+1}, z_1)$. This implies that $D_g(z_n, z_1) \leq D_g(z_{n+1}, z_1)$. This proves that $\{D_g(z_n, z_1)\}$ is a nondecreasing sequence. By combining this with the boundedness of the sequence $\{D_g(z_n, z_1)\}$, we conclude that the limit $\lim_{n \rightarrow \infty} D_g(z_n, z_1)$ exists.

Step 4. We show that $\lim_{n \rightarrow \infty} z_n = p \in \Omega$ and $\lim_{n \rightarrow \infty} \|z_{n+1} - z_n\| = 0$. Indeed, for $m > n$, we have $z_m = P_{\Omega_m}^g(z_1) \in \Omega_m \subset \Omega_n$. By choosing $y = z_m$ in (2.8), we get

$$D_g(z_m, z_n) + D_g(z_n, z_1) \leq D_g(z_m, z_1).$$

$$\text{This implies that } 0 \leq D_g(z_m, z_n) \leq D_g(z_m, z_1) - D_g(z_n, z_1). \quad (2.10)$$

Taking the limit (2.10) as $m, n \rightarrow \infty$ and using the existence of $\lim_{n \rightarrow \infty} D_g(z_n, z_1)$, we get

$$\lim_{n, m \rightarrow \infty} D_g(z_m, z_n) = 0. \quad (2.11)$$

By combining (2.11) with the boundedness of the sequence $\{z_n\}$, by Proposition 1.8, we have $\lim_{n, m \rightarrow \infty} \|z_n - z_m\| = 0$. (2.12)

This proves that $\{z_n\}$ is a Cauchy sequence in Ω . Since X is a Banach space and Ω is a closed subset of X , there exists $p \in \Omega$ such that $\lim_{n \rightarrow \infty} z_n = p$. Moreover, by choosing $m = n + 1$ in (2.11) and (2.12), we obtain $\lim_{n \rightarrow \infty} D_g(z_{n+1}, z_n) = 0$ (2.13)

$$\text{and } \lim_{n \rightarrow \infty} \|z_{n+1} - z_n\| = 0. \quad (2.14)$$

Step 5. We show that $p \in \mathcal{F}$. Indeed, since $z_{n+1} = P_{\Omega_{n+1}}^g(z_1) \in \Omega_{n+1} \subset \Omega_n$, we have

$$D_g(z_{n+1}, w_n) \leq D_g(z_{n+1}, z_n) + \gamma_n. \quad (2.15)$$

It follows from (2.9) and the boundedness of $\{D_g(z_n, z_1)\}$ that $\{D_g(u, z_n)\}$ is bounded for any $u \in \mathcal{F}$. Then, by using $\lim_{n \rightarrow \infty} \alpha_n = \lim_{n \rightarrow \infty} \beta_n = 0$, we find that $\lim_{n \rightarrow \infty} \gamma_n = 0$. (2.16)

Therefore, from (2.13), (2.15) and (2.16), we conclude that $\lim_{n \rightarrow \infty} D_g(z_{n+1}, w_n) = 0$. (2.17)

Let $u \in \mathcal{F}$. By (2.9) and the boundedness of $\{D_g(z_n, z_1)\}$, we obtain that $\{D_g(u, z_n)\}$ is bounded. By combining this with (2.7), we conclude that $\{D_g(u, w_n)\}$ is bounded. Furthermore, by Proposition 1.15, we find that g^* is bounded on bounded sets. Then ∇g^* is bounded on bounded sets. It follows from Proposition 1.10(2) that $\{w_n\}$ is bounded. By combining this with (2.17), from Proposition 1.8, we have $\lim_{n \rightarrow \infty} \|z_{n+1} - w_n\| = 0$. (2.18)

It follows from (2.14) and (2.18) that $\lim_{n \rightarrow \infty} \|z_n - w_n\| = 0$. (2.19)

Since g is uniformly Fréchet differentiable, g is uniformly continuous. Then, from (2.19) we get $\lim_{n \rightarrow \infty} \|g(z_n) - g(w_n)\| = 0$. (2.20)

Since g is uniformly Fréchet differentiable, ∇g is uniformly continuous on bounded subsets of X . Therefore, from (2.19), we have $\lim_{n \rightarrow \infty} \|\nabla g(z_n) - \nabla g(w_n)\| = 0$. (2.21)

For any $u \in \mathcal{F}$, by using similar arguments as in the proofs of (2.3) and (2.4), we obtain

$$D_g(u, u_n) \leq D_g(u, z_n) + \alpha_n \zeta(D_g(u, z_n)) + \beta_n \quad (2.22)$$

$$\text{and } D_g(u, v_n) \leq D_g(u, z_n). \quad (2.23)$$

By combining (2.22) with the boundedness of $\{D_g(u, x_n)\}$, we get that $\{D_g(u, u_n)\}$ is bounded. By Proposition 1.10(2), we get that $\{u_n\}$ is bounded. It follows from (2.23) and the boundedness of $\{D_g(u, x_n)\}$ that $\{D_g(u, v_n)\}$ is bounded. Since $\{D_g(u, v_n)\}$ is bounded and $D_g(u, S^n v_n) \leq D_g(u, v_n) + \alpha_n \zeta(D_g(u, v_n)) + \beta_n$, we find that $\{D_g(u, S^n v_n)\}$ is bounded. Thus, from Proposition 1.10(2), we get that $\{S^n v_n\}$ is bounded. Since $\{u_n\}$, $\{S^n v_n\}$ are bounded and ∇g is bounded on bounded subsets of X , we conclude that $\{\nabla g(u_n)\}$ and $\{\nabla g(S^n v_n)\}$ are bounded. Put $r = \sup_{n \in \mathbb{N}^*} \max\{\|\nabla g(u_n)\|, \|\nabla g(S^n v_n)\|\}$. Therefore,

$\nabla g(u_n), \nabla g(S^n v_n) \in B_\varepsilon = \{u \in X^* : \|u\| \leq \varepsilon\}$. By Proposition 1.14, we find that g^* is uniformly convex on bounded subsets of X^* . Therefore, by Lemma 1.16, we have

$$\begin{aligned} & g^*(b_n \nabla g(u_n) + (1 - b_n) \nabla g(S^n v_n)) \\ & \leq b_n g^*(\nabla g(u_n)) + (1 - b_n) g^*(\nabla g(S^n v_n)) - b_n (1 - b_n) \rho_\varepsilon (\|\nabla g(u_n) - \nabla g(S^n v_n)\|), \end{aligned}$$

where ρ_ε is defined as in Definition 1.11. By using Remark 1.4.(1) and the definition of V_f , we get

$$D_g(u, w_n) = D_g(u, \nabla g^*(b_n \nabla g(u_n) + (1 - b_n) \nabla g(S^n v_n))) = V_g(u, b_n \nabla g(u_n) + (1 - b_n) \nabla g(S^n v_n))$$

$$\begin{aligned}
&= g(u) - \langle b_n \nabla g(u_n) + (1 - b_n) \nabla g(S^n v_n), u \rangle + g^*(b_n \nabla g(u_n) + (1 - b_n) \nabla g(S^n v_n)) \\
&= g(u) - \langle b_n \nabla g(u_n) + (1 - b_n) \nabla g(S^n v_n), u \rangle \\
&\quad + b_n g^*(\nabla g(u_n)) + (1 - b_n) g^*(\nabla g(S^n v_n)) - b_n (1 - b_n) \rho_\varepsilon(\|\nabla g(u_n) - \nabla g(S^n v_n)\|) \\
&= b_n [g(u) - \langle \nabla g(u_n), u \rangle + g^*(\nabla g(u_n))] + (1 - b_n) [g(u) - \langle \nabla g(S^n v_n), u \rangle + g^*(\nabla g(S^n v_n))] \\
&\quad - b_n (1 - b_n) \rho_\varepsilon(\|\nabla g(u_n) - \nabla g(S^n v_n)\|) \\
&= b_n V_g(u, \nabla g(u_n)) + (1 - b_n) V_g(u, \nabla g(S^n v_n)) - b_n (1 - b_n) \rho_\varepsilon(\|\nabla g(u_n) - \nabla g(S^n v_n)\|) \\
&= b_n D_g(u, \nabla g^*(\nabla g(u_n))) + (1 - b_n) D_g(u, \nabla g^*(\nabla g(S^n v_n))) \\
&\quad - b_n (1 - b_n) \rho_\varepsilon(\|\nabla g(u_n) - \nabla g(S^n v_n)\|) \\
&= b_n D_g(u, u_n) + (1 - b_n) D_g(u, S^n v_n) - b_n (1 - b_n) \rho_\varepsilon(\|\nabla g(u_n) - \nabla g(S^n v_n)\|) \\
&\leq b_n D_g(u, u_n) + (1 - b_n) [D_g(u, v_n) + \alpha_n \zeta(D_g(u, v_n)) + \beta_n] \\
&\quad - b_n (1 - b_n) \rho_\varepsilon(\|\nabla g(u_n) - \nabla g(S^n v_n)\|). \tag{2.24}
\end{aligned}$$

Thus, by combining (2.23), (2.24) and the strictly increasing property of ζ , we get

$$\begin{aligned}
D_g(u, w_n) &\leq b_n D_g(u, u_n) + (1 - b_n) [D_g(u, z_n) + \alpha_n \zeta(D_g(u, z_n)) + \beta_n] \\
&\quad - b_n (1 - b_n) \rho_\varepsilon(\|\nabla g(u_n) - \nabla g(S^n v_n)\|). \tag{2.25}
\end{aligned}$$

By (2.22) and (2.25), we get $D_g(u, w_n) \leq D_g(u, z_n) + \gamma_n - b_n (1 - b_n) \rho_\varepsilon(\|\nabla g(u_n) - \nabla g(S^n v_n)\|)$.

This implies that $b_n (1 - b_n) \rho_\varepsilon(\|\nabla g(u_n) - \nabla g(S^n v_n)\|) \leq D_g(u, z_n) - D_g(u, w_n) + \gamma_n$. $\tag{2.26}$

Furthermore, by the property of the function D_g , we have

$$\begin{aligned}
|D_g(u, z_n) - D_g(u, w_n)| &= |-D(z_n, w_n) + \langle \nabla g(w_n) - \nabla g(z_n), u - z_n \rangle| \\
&\leq |g(z_n) - g(w_n)| + \|\nabla g(w_n)\| \cdot \|z_n - w_n\| + \|u - z_n\| \cdot \|\nabla g(w_n) - \nabla g(z_n)\|. \tag{2.27}
\end{aligned}$$

Then from (2.19), (2.20), (2.21) and (2.27), we get $\lim_{n \rightarrow \infty} |D_g(u, z_n) - D_g(u, w_n)| = 0$. $\tag{2.28}$

By (2.16), (2.26) and (2.28) that $\lim_{n \rightarrow \infty} b_n (1 - b_n) \rho_\varepsilon(\|\nabla g(u_n) - \nabla g(S^n v_n)\|) = 0$. $\tag{2.29}$

By (2.29) and $\liminf_{n \rightarrow \infty} b_n (1 - b_n) > 0$, we have $\lim_{n \rightarrow \infty} \rho_\varepsilon(\|\nabla g(u_n) - \nabla g(S^n v_n)\|) = 0$. $\tag{2.30}$

By combining (2.30) and the property of ρ_ε , we get $\lim_{n \rightarrow \infty} \|\nabla g(u_n) - \nabla g(S^n v_n)\| = 0$. $\tag{2.31}$

It follows from the assumptions of g and Proposition 1.15 that ∇g^* is uniformly continuous on bounded subsets. Thus, by (2.31), we get $\lim_{n \rightarrow \infty} \|u_n - S^n v_n\| = 0$. $\tag{2.32}$

Furthermore, by $\nabla g = (\nabla g^*)^{-1}$ and the definition of u_n , we obtain

$$\begin{aligned}
\nabla g(u_n) &= \nabla g(\nabla g^*(a_n \nabla g(z_n) + (1 - a_n) \nabla g(S^n z_n))) = a_n \nabla g(z_n) + (1 - a_n) \nabla g(S^n z_n). \text{ This} \\
&\text{leads to } \|\nabla g(u_n) - \nabla g(z_n)\| = (1 - a_n) \|\nabla g(S^n z_n) - \nabla g(z_n)\|. \tag{2.33}
\end{aligned}$$

By $\lim_{n \rightarrow \infty} a_n = 1$, the boundedness of $\{z_n\}$, (2.33), we get $\lim_{n \rightarrow \infty} \|\nabla g(u_n) - \nabla g(z_n)\| = 0$. $\tag{2.34}$

Since ∇g^* is uniformly continuous on bounded subsets, from (2.34), we have

$$\lim_{n \rightarrow \infty} \|u_n - z_n\| = 0. \quad (2.35)$$

$$\text{By combining (2.32) and (2.35), we obtain } \lim_{n \rightarrow \infty} \|z_n - S^n v_n\| = 0. \quad (2.36)$$

It follows from (2.36) and $\lim_{n \rightarrow \infty} z_n = p$ that $\lim_{n \rightarrow \infty} S^n v_n = p$. Thus, by combining this with the asymptotically regular property of S and $\|S^{n+1} v_n - p\| \leq \|S^{n+1} v_n - S^n v_n\| + \|S^n v_n - p\|$, we conclude that $\lim_{n \rightarrow \infty} S^{n+1} v_n = p$. This leads to $\lim_{n \rightarrow \infty} S(S^n v_n) = \lim_{n \rightarrow \infty} S^{n+1} v_n = p$. Since S is closed, we conclude that $Sp = p$ and hence $p \in F(S)$.

Next, we will prove that $p \in GMEP(f, \varphi, \psi)$. Since $v_n = \text{Res}_{f, \varphi, \psi}^g(z_n)$, we obtain

$$f(v_n, v) + \varphi(v) + \langle \psi(v_n), v - v_n \rangle + \langle \nabla g(v_n) - \nabla g(z_n), v - v_n \rangle \geq \varphi(v_n) \text{ for all } v \in \Omega. \quad (2.37)$$

Then, from the condition (C_2) and (2.37), we have

$$f(v, v_n) \leq -f(v_n, v) \leq \langle \psi(v_n), v - v_n \rangle + \langle \nabla g(v_n) - \nabla g(z_n), v - v_n \rangle + \varphi(v) - \varphi(v_n). \quad (2.38)$$

Furthermore, since g and ∇g are uniformly continuous on bounded subsets of X , by (2.36), we get $\lim_{n \rightarrow \infty} \|g(z_n) - g(S^n v_n)\| = \lim_{n \rightarrow \infty} \|\nabla g(z_n) - \nabla g(S^n v_n)\| = 0. \quad (2.39)$

$$\begin{aligned} \text{We have } |D_g(u, z_n) - D_g(u, S^n v_n)| &= |-D(z_n, S^n v_n) + \langle \nabla g(S^n v_n) - \nabla g(z_n), u - z_n \rangle| \\ &\leq |g(z_n) - g(S^n v_n)| + \|\nabla g(S^n v_n)\| \cdot \|z_n - S^n v_n\| + \|u - z_n\| \cdot \|\nabla g(S^n v_n) - \nabla g(z_n)\|. \end{aligned} \quad (2.40)$$

$$\text{By (2.36), (2.39) and (2.40), we get that } \lim_{n \rightarrow \infty} |D_g(u, z_n) - D_g(u, S^n v_n)| = 0. \quad (2.41)$$

For $u \in \mathcal{F}$, by Lemma 1.22 and $v_n = \text{Res}_{f, \varphi, \psi}^g(z_n)$, we find that

$$D_g(v_n, z_n) \leq D_g(u, z_n) - D_g(u, v_n) \leq D_g(u, z_n) - D_g(u, S^n v_n) + \alpha_n \zeta(D_g(u, v_n)) + \beta_n. \quad (2.42)$$

It follows from (2.41), (2.42) and $\lim_{n \rightarrow \infty} \alpha_n = \lim_{n \rightarrow \infty} \beta_n = 0$ that $\lim_{n \rightarrow \infty} D_g(v_n, z_n) = 0$. Since $\{z_n\}$ is bounded, by Proposition 1.8, we have $\lim_{n \rightarrow \infty} \|v_n - z_n\| = 0$. Since ∇g is uniformly continuous on bounded subsets, we get $\lim_{n \rightarrow \infty} \|\nabla g(z_n) - \nabla g(v_n)\| = 0$. Therefore, by using (2.38), the lower semi-continuous property of φ , the lower semi-continuous property in the second variable of f and the continuous property of ψ , we have

$$f(v, p) \leq \langle \psi(p), v - p \rangle + \varphi(v) - \varphi(p) \text{ and hence } f(v, p) + \langle \psi(p), p - v \rangle + \varphi(p) - \varphi(v) \leq 0 \text{ for all } v \in \Omega. \quad (2.43)$$

For all $t \in (0, 1]$, put $v_t = tv + (1-t)p$. Since $v, p \in \Omega$ and Ω is convex, we have $v_t \in \Omega$.

$$\text{Thus, replacing } v \text{ by } v_t \text{ in (2.43), we get } f(v_t, p) + \langle \psi(p), p - v_t \rangle + \varphi(p) - \varphi(v_t) \leq 0. \quad (2.44)$$

Then, by using the condition (C_1) , the convexity in the second variable of f , the convexity of φ and (2.44), we have

$$0 = f(v_t, v_t) = f(v_t, v_t) + \langle \psi(p), v_t - v_t \rangle + \varphi(v_t) - \varphi(v_t)$$

$$\begin{aligned}
&\leq tf(v_i, v) + (1-t)f(v_i, p) + t\langle\psi(p), v - v_i\rangle + (1-t)\langle\psi(p), p - v_i\rangle + t\varphi(v) + (1-t)\varphi(p) - \varphi(v_i) \\
&= t[f(v_i, v) + \langle\psi(p), v - v_i\rangle + \varphi(v) - \varphi(v_i)] + (1-t)[f(v_i, p) + \langle\psi(p), p - v_i\rangle + \varphi(p) - \varphi(v_i)] \\
&\leq t[f(v_i, v) + \langle\psi(p), y - v_i\rangle + \varphi(v) - \varphi(v_i)].
\end{aligned}$$

This leads to $f(v_i, v) + \langle\psi(p), v - v_i\rangle + \varphi(v) - \varphi(v_i) \geq 0$ by $t \in (0, 1]$. Letting $t \rightarrow 0^+$ and using the condition (C_3) , we have $f(p, v) + \langle\psi(p), y - p\rangle + \varphi(v) - \varphi(p) \geq 0$. This proves that $p \in GMEP(f, \varphi, \psi)$. Therefore, $p \in \mathcal{F} = F(S) \cap GMEP(f, \varphi, \psi)$.

Step 6. We show that $p = P_{\mathcal{F}}^g(z_1)$. Indeed, since $z_{n+1} = P_{\Omega_{n+1}}^g(z_1)$, by Proposition 1.9, we have

$$\langle \nabla g(z_1) - \nabla g(z_{n+1}), z_{n+1} - v \rangle \geq 0 \text{ for all } v \in \Omega_{n+1}. \text{ Let } u \in \mathcal{F}. \text{ Since } \mathcal{F} \subset \Omega_{n+1}, \text{ we get } u \in \Omega_{n+1}.$$

By choosing $v = u$ in the above inequality, we get $\langle \nabla g(z_1) - \nabla g(z_{n+1}), z_{n+1} - u \rangle \geq 0$. Taking

$n \rightarrow \infty$, using $\lim_{n \rightarrow \infty} z_n = p$ and the uniform continuous on bounded subsets of ∇g , we have

$$\langle \nabla g(z_1) - \nabla g(p), p - u \rangle \geq 0 \text{ for all } u \in \mathcal{F}. \text{ By Proposition 1.9, we find that } p = P_{\mathcal{F}}^g(z_1). \quad \square$$

Remark 2.2. (1) Theorem 2.1 is an extension of [Alizadeh & Moradlou, 2016, Theorem 3.1] from a generalized hybrid mapping in Hilbert spaces to a Bregman totally quasi-asymptotically nonexpansive mapping, and from an equilibrium problem to a generalized mixed equilibrium problem in reflexive Banach spaces.

(2) Since [Alizadeh & Moradlou, 2016, Theorem 3.1] is an extension of [Tada & Takahashi, 2007, Theorem 3.1], Theorem 2.1 is also an extension of [Tada & Takahashi, 2007, Theorem 3.1].

(3) By Remark 1.8(2), we conclude that the conclusion of Theorem 2.1 holds when S is a Bregman quasi-asymptotically nonexpansive mapping.

Finally, an example is given to illustrate for the proposed iteration.

Example 2.3. Let $X = \mathbb{R}$, $\Omega = [0, 0.9]$, $g(x) = u^2$ for all $x \in \mathbb{R}$, and $S(u) = u^2$, $\varphi(u) = 10u^2$, $\psi(u) = 2u$, $f(u, v) = -9u^2 + 4uv + 5v^2$ for all $u, v \in \Omega$. Then

(1) By calculating, we get $\nabla g(u) = 2u$, $g^*(w) = \frac{w^2}{4}$, $\nabla g^*(w) = \frac{w}{2}$ for all $u, w \in \mathbb{R}$.

(2) For all $u, v \in \mathbb{R}$, we have $D_g(u, v) = u^2 - v^2 - 2v(u - v) = (u - v)^2$.

(3) We have $F(S) = \{0\}$. Therefore, for $w \in F(S)$ and $u \in \Omega$, we obtain

$$D_g(w, S^n u) = (0 - S^n u)^2 = (u)^{2^{n+1}} \leq u^2 = D_g(0, u) = D_g(w, u).$$

This proves that S is a Bregman totally quasi-asymptotically nonexpansive mapping with $\alpha_n = \beta_n = 0$ for all $n \in \mathbb{N}^*$.

(4) By directly checking, we find that f satisfies the conditions (C_1) -(C_4).

(5) We find the formula of $w = \text{Res}_{f, \varphi, \psi}^g(u)$ for $u \in X, w \in \Omega$ as in (1.2). Indeed, $w = \text{Res}_{f, \varphi, \psi}^g(u)$

if $f(w, v) + \varphi(v) + \langle\psi(w), v - w\rangle + \langle \nabla g(w) - \nabla g(u), v - w \rangle \geq \varphi(w), v \in \Omega$. (2.45)

By substituting $f, \varphi, \psi, \nabla g$ into (2.45) and by directly calculating, we get

$$15v^2 + (8w - 2u)v + 2uw - 23w^2 \geq 0.$$

Put $h(v) = 15v^2 + (8w - 2u)v + 2uw - 23w^2$ for all $v \in \Omega$. Then $h(v)$ is a quadratic function and $\Delta = (38w - 2u)^2$. We consider the following cases.

Case 1. $\Delta > 0$. Then the equation $h(v) = 0$ has two solutions: $v_1 = w \in \Omega$ and $v_2 = \frac{-23w + 2u}{15}$. In order to $h(v) \geq 0$ for all $v \in \Omega$, we have following two cases:

Case 1.1. $v_1 = 0.9$ and $v_1 < v_2$. Then $w = v_1 = 0.9$, $v_2 = \frac{-20.7 + 2u}{15} > 0.9$, hence $u > 17.1$.

Case 1.2. $v_1 = 0$ and $v_2 < v_1$. Then $w = v_1 = 0$ and $v_2 = \frac{2u}{15} < 0$. This leads to $u < 0$.

Case 2. $\Delta \leq 0$. Then $w = \frac{u}{19}$ and $h(v) \geq 0$ for all $v \in \Omega$. Since $w \in \Omega$, we have $0 \leq \frac{u}{19} \leq 0.9$ and hence $0 \leq u \leq 17.1$. Therefore, $\text{Res}_{f,\varphi,\psi}^g(u) = w = 0$ if $u < 0$, $\text{Res}_{f,\varphi,\psi}^g(u) = w = \frac{u}{19}$ if $0 \leq u \leq 17.1$ and $\text{Res}_{f,\varphi,\psi}^g(u) = w = 0.9$ if $u > 17.1$.

By the above, all assumptions in Theorem 2.1 are satisfied with the given functions f, φ, ψ, T . Therefore, by Theorem 2.1, the sequence $\{z_n\}$ which is defined by (2.1) converges to $0 \in \mathcal{F} = F(S) \cap \text{GMEP}(f, \varphi, \psi)$. Next, by choosing $a_n = \frac{n}{n+2}$, $b_n = \frac{n+1}{3n+2}$ for all $n \in \mathbb{N}^*$, and $z_1 = 0.5 \in \Omega$, we have $P_{\mathcal{F}}^g(z_1) = \{0\}$. The sequence (2.1) becomes

$$\begin{cases} u_n = \frac{n}{n+2}z_n + \frac{2}{n+2}(z_n)^{2^n}, v_n = \frac{z_n}{19} \\ w_n = \frac{n+1}{3n+2}u_n + \frac{2n+1}{3n+2}(v_n)^{2^n}, z_{n+1} = \frac{z_n + w_n}{2}. \end{cases} \quad (2.46)$$

The convergence of iteration (2.46) is presented by the following figure.




Figure 1. The convergence of the sequence (2.46) to 0

3. Conclusion

In this paper, a hybrid iterative method is proposed for finding common elements of the solution set of a generalized mixed equilibrium problem and the fixed point set of a Bregman totally quasi-asymptotically nonexpansive mapping. After that, a strong convergence result for the proposed iteration is proved in reflexive Banach spaces. This result is an improvement of the main results in (Alizadeh & Moradlou, 2016) and (Tada & Takahashi, 2007) from a generalized hybrid mapping, a nonexpansive mapping and an equilibrium problem in Hilbert spaces to a Bregman totally quasi-asymptotically nonexpansive mapping and a generalized mixed equilibrium problem in reflexive Banach spaces. As application, we obtain the convergence result for a generalized mixed equilibrium problem and a Bregman quasi-asymptotically nonexpansive mapping in reflexive Banach spaces. Moreover, we give a numerical example to illustrate for the proposed iterative method.

❖ **Conflict of Interest:** Author have no conflict of interest to declare.

❖ **Acknowledgement:** The author was funded by Vingroup Joint Stock Company and supported by the Domestic Master/PhD Scholarship Programme of Vingroup Innovation Foundation (VINIF), Vingroup Big Data Institute (VINBIGDATA), code VNIF.2020.TS.81.

REFERENCES

- Alizadeh, S., & Moradlou, F. (2016). A strong convergence theorem for equilibrium problems and generalized hybrid mappings. *Mediterr. J. Math.*, 13(1), 379-390.
- Ambrosetti, A., & Prodi, G. (1993). *A Primer of nonlinear analysis*. Cambridge studies in Advanced Mathematics. NY: Cambridge University Press.
- Butnariu, D., & Iusem, A. N. (2000). *Totally convex functions for fixed points computation and infinite dimensional optimization*. Applied optimization, vol. 40, Kluwer Academic, NY: Dordrecht.
- Butnariu, D., & Resmerita, E. (2006). Bregman distances, totally convex functions and a method for solving operator equations in Banach spaces. *Abstr. Appl. Anal.*, 2006, 1-39.
- Censor, Y., & Lent, A. (1981). An iterative row-action method for interval convex programming. *J. Optim. Theory Appl.*, 34, 321-353.
- Chang, S. S., Wang, L., Wang, X. R., & Chan, C. K. (2014). Strong convergence theorems for Bregman totally quasi-asymptotically nonexpansive mappings in reflexive Banach spaces. *Appl. Math. Comput.*, 228, 38-48.
- Darvish, V. (2016). Strong convergence theorem for generalized mixed equilibrium problems and Bregman nonexpansive mapping in Banach spaces. *Math. Morav.*, 20(1), 69-87.
- Kumam, W., Witthayarat, U., Kumam, P., Suantai, S., & Wattanawitoon, K. (2016). Convergence theorem for equilibrium problem and Bregman strongly nonexpansive mappings in Banach spaces. *Optimization*, 65(2), 265-280.

- Kohsaka, F., & Takahashi, W. (2005). Proximal point algorithms with Bregman functions in Banach spaces. *J. Nonlinear Convex Anal.*, 6(3), 505-523.
- Naraghirad, X., & Yao, J. C. (2013). Bregman weak relatively nonexpansive mappings in Banach spaces, *Fixed Point Theory Appl.*, 2013(141), 1-43.
- Ni, R., & Wen, C. (2018). Hybrid projection methods for Bregman totally quasi-D asymptotically nonexpansive mappings. *Bull. Malays. Math. Sci. Soc.*, 41, 807-836.
- Reich, S., & S. Sabach, S. (2010). Two strong convergence theorems for a proximal method in reflexive Banach spaces. *Numer. Funct. Anal. Optim.*, 31, 22-44.
- Reich, S. & Sabach, S. (2009). A strong convergence theorem for a proximal-type algorithm in reflexive Banach spaces, *J. Nonlinear Convex Anal.*, 10, 471-485.
- Resmerita, E. (2004). On total convexity, Bregman projections and stability in Banach spaces, *J. Nonlinear Convex Anal.*, 11, 1-16.
- Sabach, S. (2011). Products of finitely many resolvents of maximal monotone mappings in reflexive Banach spaces, *SIAM J. Optim.*, 21, 1289-1308.
- Tada, A. & Takahashi, W. (2007). Weak and strong convergence theorems for a nonexpansive mapping and an equilibrium problem. *J. Optim. Theory Appl.*, 133, 359-370
- Zalinescu, Z. (2002). *Convex analysis in general vector spaces*. World Scientific, NY: River Xdge.
- Zhu, S., & Huang, J. H. (2016). Strong convergence theorems for equilibrium problem and Bregman totally quasi-asymptotically nonexpansivemapping in Banach spaces. *Acta Math. Sci. Ser. B (Xngl. Xd.)*, 36B(5), 1433-1444.

**SỰ HỘI TỤ MẠNH CỦA DÃY LẬP LẠI GHÉP
CHO BÀI TOÁN CÂN BẰNG HỖN HỢP TỔNG QUÁT
VÀ ÁNH XẠ TỰA TIỆM CẬN KHÔNG GIÃN HOÀN TOÀN BREGMAN
TRONG KHÔNG GIAN BANACH**

Nguyễn Trung Hiếu

Khoa Sư phạm Toán – Tin, Trường Đại học Đồng Tháp, Việt Nam

**Tác giả liên hệ: Nguyễn Trung Hiếu, Email: ngtrunghieu@dtu.edu.vn*

Ngày nhận bài: 20-6-2021; ngày nhận bài sửa: 20-8-2021; ngày duyệt đăng: -9-2021

TÓM TẮT

Mục đích của bài báo là kết hợp khoảng cách Bregman với phương pháp chiếu thu hẹp để giới thiệu một dãy lập lại ghép mới cho bài toán cân bằng hỗn hợp tổng quát và ánh xạ tựa tiệm cận không giãn hoàn toàn Bregman. Sau đó, với những điều kiện thích hợp, chúng tôi chứng minh rằng dãy lập lại được đề xuất hội tụ mạnh đến hình chiếu Bregman của điểm xuất phát lên giao của tập nghiệm bài toán cân bằng hỗn hợp tổng quát và tập điểm bất động của ánh xạ tựa tiệm cận không giãn hoàn toàn Bregman trong không gian Banach phản xạ. Định lý này cải tiến kết quả trong (Alizadeh & Moradlou, 2016) từ ánh xạ lai ghép tổng quát và bài toán cân bằng trong không gian Hilbert sang ánh xạ tựa tiệm cận không giãn hoàn toàn Bregman và bài toán cân bằng hỗn hợp tổng quát trong không gian Banach phản xạ. Kết quả được áp dụng cho bài toán cân bằng hỗn hợp tổng quát và ánh xạ tựa tiệm cận không giãn Bregman trong không gian Banach phản xạ. Đồng thời, một ví dụ được đưa ra để minh họa cho dãy lập lại được đề xuất.

Từ khóa: ánh xạ tựa tiếm cận không giãn hoàn toàn Bregman; bài toán cân bằng hỗn hợp tổng quát; dãy lặp lai ghép; không gian Banach phản xạ