

ANTIMICROBIAL RESISTANCE IN *Streptococcus agalactiae* IN TILAPIA (*Oreochromis* sp.) FARMING IN NORTHERN VIETNAM

Truong Thi My Hanh¹, Nguyen Thi Hanh¹, Le Thi May¹,
Truong Thi Thanh Vinh^{2*}, Dang Thi Lua¹

¹Research Institute for Aquaculture N01, Bac Ninh, Vietnam

²School of Agriculture and natural Resource, Vinh University, Nghe An, Vietnam

ARTICLE INFORMATION ABSTRACT

Journal: Vinh University
Journal of Science
Natural Science, Engineering
and Technology
p-ISSN: 3030-4563
e-ISSN: 3030-4180

Volume: 53

Issue: 2A

***Correspondence:**
thanhvinhtruong@gmail.com

Received: 04 December 2023

Accepted: 05 March 2024

Published: 20 June 2024

Citation:
Truong Thi My Hanh, Nguyen
Thi Hanh, Le Thi May, Truong
Thi Thanh Vinh, Dang Thi Lua
(2024). Antimicrobial
resistance in *Streptococcus*
agalactiae in tilapia
(*Oreochromis* sp.) farming in
Northern Vietnam.
Vinh Uni. J. Sci.
Vol. 53 (2A), pp. 15-23
doi: 10.56824/vujs.2023a157

Antimicrobial resistance (AMR) is a threat to animal and human health globally. Antimicrobials are used in aquaculture to control different bacterial diseases. However, inappropriate use of antimicrobials can accelerate the emergence of AMR. This study was conducted to determine antimicrobial resistance in *Streptococcus agalactiae*, which affects and causes diseases with high mortality in tilapia. A total of 26 *S. agalactiae* isolates were streaked from kidney/brain/liver samples of diseased farmed tilapia from 2017 to 2021 at Hai Duong, Bac Ninh, Hoa Binh, Yen Bai provinces. The sensitivity of the isolates to 7 antimicrobials was established by disc diffusion (Oxiod). The results showed that levels of resistance vary from *S. agalactiae* to DOX (34.4%), RIF (64.1%), TET (45.0%), ERY (64.8%), TMP-SMX (70%), FLO (35.3%), and AMP (75%). The resistance rate to DOX was significantly different ($p < 0.05$) compared to ERY. *S. agalactiae* has the highest resistance to 2 types of antibiotics at 19.2%, followed by resistance to 3 types (15.4%), and resistance to 6 and 7 types of antibiotics at a rate of 11.5%. The resistance in 1 type, 4 types, and 5 types of antibiotics was observed lowest at 7.7%. Vietnamese tilapia farmers need guidelines for effective antimicrobial treatment and wider take up of vaccine or sustainable farming technology to prevent infections caused by *S. agalactiae* and reduce emergence of AMR.

Keywords: Tilapia; *Streptococcus agalactiae*; antimicrobial; resistance.

OPEN ACCESS

Copyright © 2024. This is an Open Access article distributed under the terms of the [Creative Commons Attribution License \(CC BY NC\)](#), which permits non-commercially to share (copy and redistribute the material in any medium) or adapt (remix, transform, and build upon the material), provided the original work is properly cited.

1. Introduction

Antimicrobial resistance (AMR) remains a public health threat globally and south-east Asian countries have a high antimicrobial usage (AMU) and AMR [1]. Vietnam's rates of AMR are among the highest in Asia, with multi-drug-resistant infections causing thousands of deaths annually [2]. Vietnam has the highest rates of penicillin resistance (71.4%) and erythromycin resistance (92.1%) in Asia [3]. AMR is generally driven by the irrational use of antimicrobials at all levels of the health care system. However, in aquaculture and livestock production, AMU

may also contribute to AMR. Overuse of antimicrobials in livestock, use of substandard drugs, inadequate AMR surveillance, low public awareness, and inadequate regulation have all been associated with increasing AMR rates in Vietnam [4].

Fish farming is one of the fastest-growing sectors in Vietnam and plays an important role in the national economy. In 2021, the value of seafood export reached 8.9 billion US dollars; the total fishery production was 8.73 million tons, of which aquaculture production contributed 4.8 million tons [5]. The main aquaculture species cultured in Vietnam are *Pangasius* (catfish), brackish water shrimp and more recently tilapia. Oriented plan upto 2030, tilapia will become the main product farmed species after brackish water shrimp and pangasius, which will focus on developing and producing main 3 species of tilapia including: striped tilapia (*Oreochromis niloticus*); tilapia Cross-species hybrid between striped tilapia (*O. niloticus*) and green tilapia (*O. aureus*); and red tilapia (*Oreochromis* spp.) [6]. In 2020, the tilapia farming area in Vietnam reached 37,697 hectares in earth pond, over 1.2 million cubic meters (m³) in cage, strongly developed in the northern provinces such as Hai Duong, Bac Giang, Bac Ninh, Phu Tho, Hoa Binh, Hanoi, Yen Bai; Quang Ninh and some provinces in the Mekong Delta such as An Giang, Dong Thap, Hau Giang [7]. Vietnam has a tilapia production of about 270,000 tons, ranking 7th in the world after China, Indonesia; Egypt; Brazil; Philippines and Bangladesh [8], and ranked 6th in the world in terms of export volume [9].

Although the aquaculture sector, including tilapia farming, is quite established in Vietnam, it faces several challenges including diseases. A report on tilapia disease and anomalous symptoms in northern provinces (2017-2021) demonstrated that reduced feeding, scattered death, haemorrhage, cloudy/explosion eyes infected with bacteria were common and some samples also experienced parasitic disease [7]. Bacterial infections, in particular *S. agalactiae* has caused the highest mortality in Vietnamese tilapia culture across provinces [10-12]. Applying probiotics, chemicals and antimicrobials is the common practice of farmers to prevent and control disease during the grow-out phase. Approximately 70% of drugs used for animals and aquatic animals in Vietnam are antimicrobials [13]; monitoring of both livestock and aquaculture has shown that many antimicrobials critical for human health are used by farms [14]. Interviews conducted in 2022 with tilapia farmers in three northern provinces showed that amoxicillin (28.6%) and doxycycline (21.4%) were commonly used as were antimicrobials sold in pharmacies for human use (14.3%). In Bac Giang province, amoxicillin was commonly used (77.8%), whereas doxycycline/flophenicol were often used in Bac Ninh province (55.6%) [15]. However, data related to bacterial AMR in aquaculture systems are still limited, especially for tilapia pathogenic bacteria. Therefore, the determination of antibiotic resistance of bacteria isolated from infected tilapia is the aim of this research.

2. Materia and methods

2.1. Materials

Diseased tilapia samples showed clinical signs such as hemorrhage, abdominal distension, protruding eyes, and fluid in the abdominal cavity. The mortality rate of cultured fish in the pond was also recorded. Fish samples were collected from various

provinces including Hoa Binh, Hai Duong, Bac Ninh, and Yen Bai. A total of 26 strains were subjected to antibiotic analysis. The standard bacterium *E. coli* ATCC®25922™ was used as a positive control for each antibiotic tested.

The bacterial isolation method used in this study followed the protocol outlined by Buller in 2004 [16]. Liver, kidney, brain, and spleen samples from tilapia were cultured on TSA (Trypticase soy agar) nutrient medium and incubated at 28-30°C for 24-48 hours. After incubation, the bacterial cultures were subjected to proliferation culture, grams staining, and biochemical testing using API20Strep. The results of the biochemical reactions were analyzed using the Apiweb TM-API20Strep software, which identified bacterial species that matched 99% similarity with *S. agalactiae*.

The commercial discs antibiotics used in this study included doxycycline (DOX) at a concentration of 30 µg/disc, rifamycin (RIF) at 30 µl/disc, tetracycline (TET) at 30 µg/disc, erythromycin (ERY) at 15 µg/disc, trimethoprim/sulfamethoxazole (TMP-SMX) at a ratio of 1.25/23.75 µg/disc, florfenicol (FLO) at 30 µg/disc, and ampicillin (AMP) at 10 µg/disc.

2.2. Methods

The sensitivity of antibiotics was performed according to the Kirby-Bauer agar diffusion method [17]. The criteria for assessing the resistance of bacteria are based on the standards table of the Clinical and Laboratory Standards Institute (CLSI) updated in 2012.

The diameter of the zone inhibition (mm) measured using a graduated ruler after incubation at 28-30°C after 24 hours. The obtain results of resistance, sensitivity or average resistance to the tested antibiotic based on according to CLSI guidelines (Table 1).

Table 1: CLSI guidelines for the determination of antibiotic resistance in bacteria

Antibiotic paper circle	Susceptible (S)	Intermediate (I)	Resistant (R)
	Measure unit (mm)		
Doxycycline (DOX 30)	≥14	11-13	≤10
Rifamycin (RIF 30)	≥20	17-19	≤16
Tetracycline (TET 30)	≥15	12-14	≤11
Erythromycine (ERY 15)	≥23	14-22	≤13
Trimethoprim/sulfamethoxazole (TMP-SMX 1.25/23.75)	≥17	13-16	≤12
Florfenicol (FLO 30)	≥20	17-19	≤16
Ampiciline (AMP 10)	≥17	14-16	≤13

Excel 2010 software is used for descriptive statistical analysis and the differences between formulas were investigated using LSD on SPSS 20 software.

3. Results and discussion

3.1 Current status of resistance to some antibiotics of *Streptococcus agalactiae*

The analysis results indicated that *S. agalactiae* strains isolated from diseased tilapia samples exhibited varying rates of antibiotic resistance, ranging from 34.4% to

75% for the seven antibiotics investigated (Table 2). It is worth noting that the resistance rate for the antibiotic doxycycline (DOX) was 34.4%, which was significantly different ($p < 0.05$) from the resistance rate observed for Erythromycin (ERY). These findings align with previous studies conducted by Dang Thi Lua and Truong Thi My Hanh in 2019, which demonstrated that *Streptococcus* sp. isolated from tilapia and climbing perch cultivated in Phu Tho and Hai Duong exhibited resistance to AMP and TMP-SMX [18]. Additionally, another study conducted by Phuoc et al. in 2015 found that *Streptococcus* sp. causing disease in tilapia cultivated in the Mekong Delta showed resistance to various antibiotics, including ofloxacin, nitrofurantoin, tetracycline, oxacillin, bacitracin, ampicillin, amoxicillin, and streptomycin [19]. Thereby, the antibiotic resistance of *Streptococcus* sp. isolated from cultured tilapia has been clearly identified. However, there are differences in the types of antibiotics used in different research areas. This is because each region of aquaculture uses different types of medications.

Some investigation results on antibiotic use in different aquaculture areas demonstrated: In An Giang province, sulfonamide and trimethoprim were mostly used (95.0%), followed by amoxicillin (55.0%), florfenicol (50.0%). In Tien Giang province, there were 8 antibiotics, among those the most used were sulfonamide and trimethoprim (87.0%), followed by amoxicillin (56.5%). In Dong Thap province, there were 11 antibiotics included amoxicillin (86.4%), sulfonamide and trimethoprim (81.8%), ciprofloxacin (63.6%), oxytetracycline (59.1%), florfenicol (59.1%) and enrofloxacin (50.0%). The tilapia farmers in Vinh Long province have used 6 types of antibiotics in the treatment of fish diseases. The most used products are still sulfonamide and trimethoprim (81.0%), amoxicillin (76.2%), florfenicol (76.2%), doxycycline (71.4%) and other antibiotics such as ampicillin, colistin, gentamycin, florfenicol and doxycycline, erythromycin, tobramycin, etc. with low usage rates [12]. In Hai Duong province, the usage rate of amoxicillin is highest (28.6%), followed by doxycycline (21.4%), especially sometimes using human medicine for fish (14.3%). Similar to Hai Duong, 77.8% of households in Bac Giang use amoxicillin, then trimethoprim-sulfamethoxazole (66.7%), doxycycline (44.4%), while tilapia farmers in Bac Ninh commonly use a combination of two antibiotics, doxycycline and florfenicol (55.6%) [15].

In terms of sensitivity, the sensitivity rates of the 7 selected antibiotics for research on *S. agalactiae* ranged from 12.5% to 61.2%. The highest sensitivity rate of DOX at 61.2% demonstrated a significant difference ($p < 0.05$) compared to AMP (12.5%), while the other 5 antibiotics did not show any significant difference ($p > 0.05$) among themselves as well as compared to other experimental antibiotics (Table 2). Meanwhile, the intermediate results of 7 drugs tested for *S. agalactiae* ranged from 4.4-29.5% and there was no significant difference between them (Table 2). The TET and DOX antibiotics of the tetracycline group have high sensitivity to *Streptococcus* sp. has also been report in tilapia farming areas in Phu Tho and Hai Duong [18].

Table 2: Rate of antimicrobial resistance of *S. agalactiae* to some antibiotics

No.	Antibiotic paper circle	Resistant (R) (%)	Intermediate (I) (%)	Susceptible (S) (%)
1	Doxycycline (DOX 30)	34.4±10.2 ^a	4.4±4.4 ^a	61.2±15.8 ^b

No.	Antibiotic paper circle	Resistant (R) (%)	Intermediate (I) (%)	Susceptible (S) (%)
2	Rifamycin (RIF 30)	61.4±18.6 ^{ab}	5.7±5.7 ^a	32.9±19.1 ^{ab}
3	Tetracycline (TET 30)	45.0±17.8 ^{ab}	25.0±13.1 ^a	30.0±17.9 ^{ab}
4	Erythromycine (ERY 15)	64.8±17.0 ^b	25.2±8.0 ^a	30.0±20.0 ^{ab}
5	Trimethoprim/sulfamethoxazole (TMP-SMX 1.25/23.75)	70.0±20 ^{ab}	10.0±10.0 ^a	20.0±20.0 ^{ab}
6	Florfenicol (FLO 30)	35.3±19.4 ^{ab}	29.5±18.7 ^a	35.2±18.4 ^{ab}
7	Ampiciline (AMP 10)	75.0±14.4 ^{ab}	12.5±12.5 ^a	12.5±12.5 ^a

Note: The data in the table are the Mean ± Standard error. The data in the same column containing different letters are statistically significant ($p < 0.05$)

Furthermore, the analysis of the inhibition diameter of *S. agalactiae* samples collected from 2017 to 2021 for the same antibiotic revealed a gradual decrease in the sterile ring diameters over time, as indicated in Figure 2. This observation further supports the existence of antibiotic resistance in *S. agalactiae*. In 2017, the average inhibited ring diameter for the tested antibiotics ranged from 15-30 mm. However, in subsequent years, there was a sharp decline in the average inhibited ring diameter, particularly for TMP-SMX, where the average inhibited ring diameter was only 2.5mm in 2021. The average inhibited ring diameter of *S. agalactiae* for DOX, RIF, TET, and ERY also exhibited a gradual decrease over time from 2017 to 2021, with significant reductions observed during the periods of 2017-2018 and 2020-2021, resulting in a sterile diameter average reduction of 10-15 mm (Figure 1).




Figure 1: The average inhibition diameter of *S. agalactiae* with some antibiotics

3.2 Multi-antibiotic resistance of *Streptococcus agalactiae*

The study results indicate that *S. agalactiae* is both mono and multi-antibiotic resistant. *S. agalactiae* has the highest rate of resistance to 2 types of antibiotics at 19.2%, followed by resistance to 3 types of antibiotics (15.4%), and finally resistance to 6 and 7 types of antibiotics at a rate of 11.5%. The lowest resistance was seen in 1 type, 4 types, and 5 types of antibiotics at 7.7% (Figure 2). Multi-drug resistance in pathogenic bacteria in aquaculture has been studied and identified in various different farming subjects. *Streptococcus* sp. isolated from samples of carp and catfish farming in Phu Tho and Hai Duong provinces showed resistance to at least 2 antibiotics and up to 6 antibiotics [18]. *Vibrio* sp. obtained from shrimp in Long An, Ben Tre, Bac Lieu, and Ninh Thuan provinces exhibited 100% strain resistance to 1 antibiotic, 95% strain resistance to 4 antibiotics, with over 50% of bacterial strains showing resistance to more than 10 types of antibiotics, including one strain resistant to 21 types of investigated antibiotics [20]. *V. parahaemolyticus*, the causative agent of AHPND in farmed shrimp in Quynh Luu district, Nghe An province, demonstrated the highest rate of resistance to 4 types of antibiotics (33.3%), followed by resistance to 6 and 3 types of drugs (22.2%), and finally resistance to 2 and 5 types of drugs (11.1%) [21]. *Edwardsiella ictaluri*, which causes disease in farmed Tra catfish in the Mekong Delta, was found to have up to 86% of strains showing resistance to 3 or more drugs, and 70% of strains resistant from 6 to 9 types of drugs [22]. *Listeria seeligeri* and *L. welshimeri* bacteria showed resistance to 2 types of antibiotics at a rate of 10.9% [23]. Therefore, the phenomenon of multi-drug resistance in bacteria has been identified in aquaculture, posing a potential risk of transferring multi-drug resistance genes from pathogenic bacteria in aquatic animals (such as shrimp) to bacteria causing diseases in humans and livestock.




Figure 2: Mono and multi-antibiotic resistance of *S. agalactiae*

4. Conclusion

S. agalactiae, isolated from diseased tilapia samples in Hai Duong, Bac Ninh, Hoa Binh, and Yen Bai provinces, exhibited varying resistance rates to different antibiotics. The resistance rates were as follows: DOX (34.4%), RIF (64.1%), TET (45.0%), ERY (64.8%), TMP-SMX (70%), FLO (35.3%), and AMP (75%). The resistance rate to DOX was significantly different ($p < 0.05$) compared to ERY.

Among the resistant strains of *S. agalactiae*, the highest rate was observed for resistance to 2 types of antibiotics (19.2%). This was followed by resistance to 3 types of antibiotics (15.4%), then resistance to both 6 and 7 antibiotics at a rate of 11.5%. On the other hand, the lowest resistance rates were seen in strains resistant to only 1 antibiotic, 4 antibiotics, and 5 antibiotics, at 7.7%.

5. Proposal

In tilapia farming areas, including Hai Duong, Bac Ninh, Hoa Binh, and Yen Bai, it has been indicated that *S. agalactiae* exhibits resistance to certain antibiotics. Therefore, before choosing for antibiotic treatment for fish diseases, it is essential to collect samples and assess the disc antibiotic diffusion test.

REFERENCES

- [1] Zellweger R. M., Carrique-Mas J., Limmathurotsakul D. et al., "A current perspective on antimicrobial resistance in Southeast Asia," *J. Antimicrob Chemother*, 72: 2963, 2017. DOI: 10.1093/jac/dkx260
- [2] Division of Global Health Protection, Global Health, Centers for Disease Control and Prevention (DGHP, GH, CDC), "Vietnam Tracks Multi-drug Resistant Bacteria," 2018. Available: <https://www.cdc.gov/globalhealth/healthprotection/fieldupdates/winter-2017/vietnam-tracks-bacteria.html>
- [3] Song J. H. et al., "High prevalence of antimicrobial resistance among clinical *Streptococcus pneumoniae* isolates in Asia (an ANSORP study)," *Antimicrob Agents Chemother*, 48(6): p. 2101, 2004. DOI: 10.1128/AAC.48.6.2101-2107.2004
- [4] Kinh N. V. et al., "Antibiotic use and resistance in emerging economies: a situation analysis for Viet Nam," *BMC Public Health*, 13:1158, 2013. DOI: 10.1186/1471-2458-13-1158
- [5] Dfish annual report, "Directorate of Fisheries," Ministry of Agriculture and Rural Development, Vietnam, 2021.
- [6] Ministry of Agriculture and Rural Development, *Decision 1639/QĐ-BNN-TCTS, Regarding approval of plan for development of tilapia fish farming to 2020, orientation towards 2030*, Hanoi, 2016.
- [7] Hanh. M. H. T, T. H. Nguyen, M. Q. Nguyen, T. M. Le, T. N. Nguyen, T. B. Phan, T. T. V. Truong and T. V. Phan, "Cause of abnormal in tilapia (*Oreochromis* sp.) in some northern province 2017-2021," *Journal of Agriculture and Rural Development*, vol. 10, pp. 154-159, 2022.

- [8] FAO, *The international technical seminar on tilapia health on 1-3/12/2021*, 2020.
- [9] VASEP, *The export market and value export of Tilapia in Vietnam and List of enterprises exporting tilapia in Vietnam from 2018 to 2020*, 2021.
- [10] Nghia N. T. and Oanh D. T. H., "Pathogenicity of *Edwardsiella ictaluri* on red tilapia (*Oreochromis* sp.)," *Sciences Journal of Can Tho University*, 55(3b), pp. 123-131, 2019. DOI: 10.22144/ctu.jvn.2019.084
- [11] Hanh T. T. M., Nhat P. T. H, Minh L. H, Nghia N. H., Vinh T. T. T. and Van, P. T., "Some main characteristics of *Streptococcus agalactiae* cause disease on tilapia (*Oreochromis* sp.) in brackish water environment," *Journal of Agriculture and Rural Development*, pp. 73-79, 2020.
- [12] T. M. Phu et al., "The use of drug, chemical, and probiotic in red tilapia (*Oreochromis* sp.) cage culture in Mekong Delta, Vietnam," *Can Tho University Journal of Sciences*: 51b, pp. 80-87. DOI: 10.22144/ctu.jvn.2017.082
- [13] N. V. Kinh, *Situation analysis: Antibiotic use and resistance in Viet Nam*, GARP - Viet Nam Working Group, 2010.
- [14] Vietnam Ministry of Health, *Vietnam National Drug Formulary, 2nd edn.*, Hanoi: Vietnam medical publisher, 2009.
- [15] Hanh T. T. M., Nguyen T. N., Le T. M., Truong T. T. V. and Phan T. V., "Current disease and antimicrobial used in tilapia (*Oreochromis* sp.) cultured in Hai Duong, Bac Ninh and Bac Giang," *TNU Journal of Science and Technology* 228(05): 410-414, 2023. DOI: 10.34238/tnu-jst.7476
- [16] Buller N. B., *Bacteria from Fish and Other Aquatic Animals: A Practical Identification Manual*, CABI Publishing, 2004. DOI: 10.1079/9780851997384.0000
- [17] Bauer A. W., Kirby M. D., Sherris J. C. and Turck, M., (1966). Antibiotic Susceptibility Testing by a Standardized Single Disk Method. *American Journal of Clinical Patholog.* Volume 45, April 1966, Pages 493-496. DOI: 10.1093/ajcp/45.4_ts.493
- [18] Lua D. T and Hanh T. T. M., "Determination of antibiotic resistance of *Streptococcus* spp causing disease in tilapia and square-headed perch cultured in some northern provinces," *Journal of Veterinary Science and Technology*, volume 26, issue 2, pp. 26-33, 2019.
- [19] Phuoc N. N., Luu T. N. H, Nguyen T. S., Nguyen D. Q. A., Truong T. H. and Le V. B. D., "Study on biochemical characteristics of *Streptococcus* sp. pathogens in tilapia cultured in the Mekong Delta, Vietnam," *Hue University Journal of Science: Agriculture and Rural Development*, vol. 104, no. 5, 2015. DOI:10.26459/jard.v104i5.2963
- [20] Ho K. D., Truyen N. D. H and Luu T. T. T., "Antibiotic resistance of *Vibrio* spp. on shrimp (*Litopenaeus vannamei*)," *Journal of Fisheries Science and Technology Nha Trang University*, no. 4, pp. 26-32, 2019.
- [21] Hanh T. T. M., Phạm T. Y, Huynh T. M. L., Phan T. V., Nguyen D. V. and Truong T. T. V., "Current status of antibiotic usages and resistance of *Vibrio parahaemolyticus* causing acute hepatopancreatic necrosis disease on shrimp in

- Quynh Luu, Nghe An,” *Journal of Fisheries Science and Technology Nha Trang University*, no. 4, page 57, 2016.
- [22] Dung T. T. D. et al., “The phenomenon of multi-resistance on bacteria *Edwardsiella ictaluri* causing pyogenic liver disease in catfish (*Pangasianodon hypophthalmus*) in Mekong Delta,” *Journal of Sciences Can Tho University*, pp. 219-227, 2010.
- [23] Vengadesh L., Wai F. Yin, Learn-H. Lee and K. G. C., “Prevalence and antimicrobial susceptibility of *Vibrio parahaemolyticus* isolated from retail shrimps in Malaysia,” *Front. Microbiol.*, 30/01/2015. DOI: 10.3389/fmicb.2015.00033.

TÓM TẮT

KHÁNG KHÁNG SINH CỦA *Streptococcus agalactiae* GÂY BỆNH TRÊN CÁ RÔ PHI (*Oreochromis sp.*) Ở MIỀN BẮC VIỆT NAM

Trương Thị Mỹ Hạnh¹, Nguyễn Thị Hạnh¹, Lê Thị Mây¹,
Trương Thị Thành Vinh², Đặng Thị Lua¹

¹Viện nghiên cứu Nuôi trồng thủy sản 1, Bắc Ninh, Việt Nam

²Viện Nông Nghiệp và Tài Nguyên, Đại học Vinh, Nghệ An, Việt Nam

Ngày nhận bài 04/12/2023, ngày nhận đăng 05/3/2024

Kháng kháng sinh là mối đe dọa đối với sức khỏe động vật và con người trên toàn cầu. Thuốc kháng sinh được sử dụng trong nuôi trồng thủy sản để trị các bệnh do vi khuẩn. Nhưng việc sử dụng thuốc kháng sinh không phù hợp có thể dẫn đến kháng kháng sinh. Nghiên cứu này được thực hiện nhằm xác định tình trạng kháng kháng sinh của vi khuẩn *Streptococcus agalactiae*. Với tổng số 26 *S. agalactiae* được phân lập từ các mẫu thận/não/gan của cá rô phi nuôi bị bệnh từ năm 2017 đến 2021 tại các tỉnh Hải Dương, Bắc Ninh, Hòa Bình, Yên Bái. Độ nhạy của vi khuẩn phân lập với 7 loại kháng sinh được xác định bằng phương pháp khuếch tán đĩa (Oxiod). Kết quả cho thấy mức độ kháng khác nhau, với DOX (34,4%), RIF (64,1%), TET (45,0%), ERY (64,8%), TMP-SMX (70%), FLO (35,3%) và AMP (75%). Tỷ lệ *S. agalactiae* kháng với DOX khác biệt rõ rệt ($p < 0,05$) với ERY ($p < 0,05$). *S. agalactiae* cho kết quả kháng cao nhất với 2 loại kháng sinh (19,2%), tiếp đến kháng với 3 loại kháng sinh (15,4%) và kháng với 6 và 7 loại kháng sinh chiếm 11,5%. Tỷ lệ kháng với 1, 4 và 5 loại kháng sinh là thấp nhất chiếm tỷ lệ 7,7%. Người nuôi cá rô phi Việt Nam cần có hướng dẫn về điều trị kháng sinh hiệu quả và sử dụng vắc-xin hoặc công nghệ nuôi bền vững rộng rãi hơn để ngăn ngừa nhiễm khuẩn do *S. agalactiae* gây ra và giảm sự xuất hiện của kháng kháng sinh.

Từ khóa: Cá rô phi; vi khuẩn *Streptococcus agalactiae*; kháng sinh; kháng kháng sinh.