

ILL-POSEDNESS AND STABILITY ESTIMATE FOR THE HEAT EQUATION BACKWARD IN TIME WITH DIRICHLET AND INTEGRAL BOUNDARY CONDITIONS

Nguyen Van Duc^{1,*}, Phan Hoai Linh²

¹Department of Mathematics, Vinh University, Nghe An, Vietnam

²Master's student course 30, Analytical Mathematics,
Department of Mathematics, Vinh University, Nghe An, Vietnam

ARTICLE INFORMATION ABSTRACT

Journal: Vinh University
Journal of Science
Natural Science, Engineering
and Technology
p-ISSN: 3030-4563
e-ISSN: 3030-4180

Volume: 53

Issue: 4A

***Correspondence:**

ducnv@vinhuni.edu.vn

Received: 05 August 2024

Accepted: 17 October 2024

Published: 20 December 2024

Citation:

Nguyen Van Duc, Phan Hoai
Linh (2024). Ill-posedness and
stability estimate for the heat
equation backward in time with
Dirichlet and integral boundary
conditions. *Vinh Uni. J. Sci.*
Vol. 53 (4A), pp. 90-98
doi: 10.56824/vujs.2024a079a

OPEN ACCESS

Copyright © 2024. This is an
Open Access article distributed
under the terms of the Creative
Commons Attribution License
(CC BY NC), which permits non-
commercially to share (copy and
redistribute the material in any
medium) or adapt (remix,
transform, and build upon the
material), provided the original
work is properly cited.

In this paper, we first prove that the heat equation backward in time with Dirichlet and integral boundary conditions is an ill-posed problem. Then, we establish a stability estimate of Hölder type for this ill-posed problem.

Keywords: Heat equation backward; ill-posed problem; stability estimate.

1. Introduction

Let l and T be positive real numbers. Consider the problem of determining $u(x, t)$ satisfying

$$\begin{cases} u_l - u_{xx} = 0, 0 < x < l, 0 < t < T \\ u(0, t) = 0, \int_0^l u(x, t) dx = 0, 0 \leq t \leq T \end{cases} \quad (1)$$

with measurement data at $t = T$:

$$u(x, T) = \varphi(x), x \in [0, l] \quad (2)$$

where φ is a given function.

Problem (1)-(2) is ill-posed in the sense of Hadamard (see Theorem 2.1). Therefore, stability estimates and regularization methods are desired.

In recent years, there have been many research papers on parabolic equations with integration conditions ([1], [2], [3], [4], [5], [6], [7]). In paper [1], we have established a stability estimate result of the Hölder type for heat equation backward in time with Neumann and integral boundary conditions. However, the technique in paper [1] cannot be used to establish stability estimate results for heat equation backward in time with Dirichlet and integral boundary conditions because the two boundary conditions, Dirichlet and Neumann, are different. Furthermore, to our knowledge, there is so far not any result on stability estimates of heat equation backward in time with Dirichlet and integral boundary conditions (1)-(2). This motivates us to establish a stability estimate result for problem (1)-(2).

The purpose of this paper is to prove that problem (1)-(2) is an ill-posed problem and propose a stability estimate result of the Hölder type for this problem. We present these results in Section 2.

2 Main results

For simplicity of notation, in this section we denote $\| \cdot \|_{L^2(0,l)}$ by $\| \cdot \|$.

Theorem 2.1. (ill-posedness) *Problem (1)-(2) is an ill-posed problem.*

Proof. Set

$$\begin{aligned} u^n(x, t) &= \frac{1}{n} e^{\left(\frac{2\pi n}{l}\right)^2 (T-t)} \sin \frac{2\pi n x}{l}, \quad 0 \leq x \leq l, \quad 0 \leq t \leq T, \\ \varphi^n(x) &= \frac{1}{n} \sin \frac{2\pi n x}{l}, \quad 0 \leq x \leq l, \\ u(x, t) &= 0, \quad 0 \leq x \leq l, \quad 0 \leq t \leq T, \\ \varphi(x) &= 0, \quad 0 \leq x \leq l. \end{aligned}$$

It follows that

$$\begin{aligned} u^n(0, t) &= 0, \quad 0 \leq t \leq T, \\ u_t^n(x, t) &= - \left(\frac{2\pi n}{l}\right)^2 \frac{1}{n} e^{\left(\frac{2\pi n}{l}\right)^2 (T-t)} \sin \frac{2\pi n x}{l}, \quad 0 \leq x \leq l, \quad 0 \leq t \leq T, \\ u_{xx}^n(x, t) &= \frac{2\pi n}{l} \frac{1}{n} e^{\left(\frac{2\pi n}{l}\right)^2 (T-t)} \cos \frac{2\pi n x}{l}, \quad 0 \leq x \leq l, \quad 0 \leq t \leq T, \\ u_{xx}^n(x, t) &= - \left(\frac{2\pi n}{l}\right)^2 \frac{1}{n} e^{\left(\frac{2\pi n}{l}\right)^2 (T-t)} \sin \frac{2\pi n x}{l}, \quad 0 \leq x \leq l, \quad 0 \leq t \leq T, \end{aligned}$$

$$\begin{aligned} \int_0^l u^n(x, t) dx &= \int_0^l \frac{1}{n} e^{\left(\frac{2\pi n}{l}\right)^2 (T-t)} \sin \frac{2\pi n x}{l} dx \\ &= \frac{1}{n} e^{\left(\frac{2\pi n}{l}\right)^2 (T-t)} \int_0^l \sin \frac{2\pi n x}{l} dx \\ &= -\frac{1}{n} e^{\left(\frac{2\pi n}{l}\right)^2 (T-t)} \frac{l}{2\pi n} \cos \frac{2\pi n x}{l} \Big|_0^l \\ &= -\frac{l}{n} e^{\left(\frac{2\pi n}{l}\right)^2 (T-t)} \frac{\cos 2\pi n - \cos 0}{2\pi n} = 0, \quad 0 \leq t \leq T, \\ u^n(x, T) &= \frac{1}{n} \sin \frac{2\pi n x}{l} = \varphi^n(x), \quad 0 \leq x \leq l, \end{aligned}$$

$$\begin{aligned} u_t(x, t) = 0 &= u_{xx}(x, t), \quad 0 \leq x \leq l, \quad 0 \leq t \leq T, \\ u(0, t) = 0, \quad \int_0^l u(x, t) dx &= 0, \quad 0 \leq t \leq T, \\ u(x, T) = 0 &= \varphi(x), \quad 0 \leq x \leq l. \end{aligned}$$

Therefore $u^n(x, t)$ satisfies the following conditions

$$\begin{cases} u_t^n - u_{xx}^n = 0, & 0 < x < l, 0 < t < T \\ u^n(0, t) = 0, \quad \int_0^l u^n(x, t) dx = 0, & 0 \leq t \leq T \\ u^n(x, T) = \varphi^n(x), & 0 \leq x \leq l \end{cases}$$

and $u(x, t)$ satisfies the following conditions

$$\begin{cases} u_t - u_{xx} = 0, & 0 < x < l, 0 < t < T \\ u(0, t) = 0, \quad \int_0^l u(x, t) dx = 0, & 0 \leq t \leq T \\ u(x, T) = \varphi(x), & 0 \leq x \leq l. \end{cases}$$

On the other hand, we have

$$\begin{aligned} \|\varphi^n - \varphi\| &= \left(\int_0^l (\varphi^n(x))^2 dx \right)^{\frac{1}{2}} = \left(\int_0^l \left(\frac{1}{n} \sin \frac{2\pi nx}{l} \right)^2 dx \right)^{\frac{1}{2}} \\ &= \frac{1}{n} \left(\int_0^l \sin^2 \frac{2\pi nx}{l} dx \right)^{\frac{1}{2}} = \frac{1}{n} \left(\int_0^l \frac{1 - \cos \frac{4\pi nx}{l}}{2} dx \right)^{\frac{1}{2}} \\ &= \frac{1}{n} \left(\frac{1}{2} x \Big|_0^l - \frac{l \sin \frac{4\pi nx}{l}}{8\pi n} \Big|_0^l \right)^{\frac{1}{2}} \\ &= \frac{\sqrt{l}}{\sqrt{2n}} \rightarrow 0 \text{ as } n \rightarrow +\infty. \end{aligned}$$

However, for $0 \leq t < T$ we obtain

$$\begin{aligned} \|u^n(\cdot, t) - u(\cdot, t)\| &= \left(\int_0^l u^n(x, t)^2 dx \right)^{\frac{1}{2}} = \left(\int_0^l \left(\frac{1}{n} e^{\left(\frac{2\pi n}{l}\right)^2 (T-t)} \sin \frac{2\pi nx}{l} \right)^2 dx \right)^{\frac{1}{2}} \\ &= \frac{1}{n} e^{\left(\frac{2\pi n}{l}\right)^2 (T-t)} \left(\int_0^l \left(\sin \frac{2\pi nx}{l} \right)^2 dx \right)^{\frac{1}{2}} \\ &= \frac{\sqrt{l}}{\sqrt{2n}} e^{\left(\frac{2\pi n}{l}\right)^2 (T-t)} \rightarrow +\infty \text{ as } n \rightarrow +\infty. \end{aligned}$$

This proves that problem (1)-(2) is an ill-posed problem.

Hence, the theorem is proved. □

Theorem 2.2. (Stability estimate) Let $u^1(x, t)$ and $u^2(x, t)$ be solutions of problem (1). Assume that there exists a constant $C \leq 0$ such that

$$u^i(l, t) = Cu_x^i(0, t), i = 1, 2. \tag{3}$$

If $u^1(x, t)$ and $u^2(x, t)$ satisfy

$$\|u^i(\cdot, 0)\| \leq E, i = 1, 2. \tag{4}$$

and $\|u^1(\cdot, T) - u^2(\cdot, T)\| \leq \delta$ where δ and E are some positive constants then

$$\|u^1(\cdot, t) - u^2(\cdot, t)\| \leq 2^{(1-\frac{t}{T})} \delta^{\frac{t}{T}} E^{1-\frac{t}{T}}, \forall t \in [0, T].$$

Proof. Set

$$u(x, t) = u^1(x, t) - u^2(x, t), (x, t) \in [0, l] \times [0, T].$$

We have

$$\begin{aligned} u_t(x, t) &= u_t^1(x, t) - u_t^2(x, t) \\ &= u_{xx}^1(x, t) - u_{xx}^2(x, t) \\ &= u_{xx}(x, t), 0 < x < l, 0 < t < T \\ u(0, t) &= u^1(0, t) - u^2(0, t) = 0 \\ \int_0^l u(x, t) dx &= \int_0^l (u^1(x, t) - u^2(x, t)) dx = 0, 0 \leq t \leq T. \end{aligned}$$

Thus, the function $u(x, t)$ satisfies the following conditions

$$\begin{cases} u_t - u_{xx} = 0, 0 < x < l, 0 < t < T \\ u(0, t) = 0, \int_0^l u(x, t) dx = 0, 0 \leq t \leq T. \end{cases} \tag{5}$$

Set $h(t) = \int_0^l (u(x, t))^2 dx = \|u(\cdot, t)\|^2, t \in [0, T]$. By the integration by part, we have

$$\begin{aligned} h'(t) &= 2 \int_0^l u(x, t) u_t(x, t) dx \\ &= 2 \int_0^l u(x, t) u_{xx}(x, t) dx \\ &= 2 \int_0^l u(x, t) d(u_x(x, t)) = 2u(x, t) u_x(x, t) \Big|_0^l - 2 \int_0^l (u_x(x, t))^2 dx \\ &= 2u(l, t) u_x(l, t) - 2u(0, t) u_x(0, t) - 2 \int_0^l (u_x(x, t))^2 dx \\ &= 2u(l, t) u_x(l, t) - 2 \int_0^l (u_x(x, t))^2 dx \text{ (since } u(0, t) = 0) \end{aligned}$$

$$\begin{aligned}
 h''(t) &= 2u_t(l, t)u_x(l, t) + 2u(l, t)u_{xt}(l, t) - 4 \int_0^l u_x(x, t)u_{xt}(x, t)dx \\
 &= 2u_t(l, t)u_x(l, t) + 2u(l, t)u_{xt}(l, t) - 4u_x(x, t)u_t(x, t) \Big|_0^l + 4 \int_0^l (u_t(x, t))^2 dx \\
 &= 2u_t(l, t)u_x(l, t) + 2u(l, t)u_{xt}(l, t) - 4u_x(l, t)u_t(l, t) \\
 &\quad + 4 \int_0^l (u_t(x, t))^2 dx, \text{ (since } u(0, t) = 0, u_t(0, t) = 0) \\
 &= 2u(l, t)u_{xt}(l, t) - 2u_x(l, t)u_t(l, t) + 4 \int_0^l (u_t(x, t))^2 dx. \tag{6}
 \end{aligned}$$

From the assumption

$$u^i(l, t) = Cu_x^i(0, t), i = 1, 2 \tag{7}$$

we have

$$\begin{cases} u^1(l, t) = Cu_x^1(0, t) \\ u^2(l, t) = Cu_x^2(0, t). \end{cases} \tag{8}$$

This implies that

$$Cu_{xx}(0, t) = Cu_x^1(0, t) - Cu_x^2(0, t) = u^1(l, t) - u^2(l, t) = u(l, t). \tag{9}$$

It follows from

$$\int_0^l u(x, t)dx = 0$$

that

$$\int_0^l u_t(x, t)dx = 0.$$

This implies that

$$\int_0^l u_{xx}(x, t)dx = 0 \text{ (since } u_t(x, t) = u_{xx}(x, t)).$$

Therefore, we obtain

$$u_x(l, t) = u_x(0, t). \tag{10}$$

From (9) and (10), we have

$$u(l, t) = Cu_x(l, t).$$

This implies that

$$\begin{aligned} h'(t) &= 2u(l, t)u_x(l, t) - 2 \int_0^l (u_x(x, t))^2 dx \\ &= 2Cu_x^2(l, t) - 2 \int_0^l (u_x(x, t))^2 dx \leq 0 \text{ (since } C \leq 0), \forall t \in (0, T). \end{aligned}$$

If $C < 0$, then from the above equality, we have

$$\begin{aligned} 2u(l, t)u_{xt}(l, t) - 2u_x(l, t)u_t(l, t) \\ &= \frac{2}{C}u(l, t)u_t(l, t) - \frac{2}{C}u(l, t)u_t(l, t) \\ &= 0. \end{aligned} \tag{11}$$

If $C = 0$ then $u(l, t) = 0$. This implies that $u_t(l, t) = 0$. Therefore, we get

$$2u(l, t)u_{xt}(l, t) - 2u_x(l, t)u_t(l, t) = 0. \tag{12}$$

From (6), (11) and (12), we have

$$h''(t) = 4 \int_0^l (u_t(x, t))^2 dx. \tag{13}$$

Due to the Cauchy-Schwarz inequality, we have

$$\left(\int_0^l u(x, t)u_t(x, t) dx \right)^2 \leq \int_0^l (u(x, t))^2 dx \int_0^l (u_t(x, t))^2 dx.$$

This implies that

$$\begin{aligned} h(t)h''(t) - (h'(t))^2 \\ &= 4 \left(\int_0^l (u(x, t))^2 dx \int_0^l (u_t(x, t))^2 dx - \left(\int_0^l u(x, t)u_t(x, t) dx \right)^2 \right) \\ &\geq 0, \forall t \in [0, T]. \end{aligned}$$

This proves that f is a logarithmically convex function. Therefore, we have

$$h(t) \leq h(T)^{\frac{t}{T}} h(0)^{1-\frac{t}{T}}, \forall t \in [0, T]$$

or

$$\|u(\cdot, t)\| \leq \|u(\cdot, T)\|^{\frac{t}{T}} \|u(\cdot, 0)\|^{1-\frac{t}{T}}, \forall t \in [0, T]. \tag{14}$$

On the other hand, we have

$$\|u(\cdot, T)\| = \|u^1(\cdot, T) - u^2(\cdot, T)\| \leq \delta \tag{15}$$

and

$$\|u(\cdot, 0)\| = \|u^1(\cdot, 0) - u^2(\cdot, 0)\| \leq \|u^1(\cdot, 0)\| + \|u^2(\cdot, 0)\| \leq E + E = 2E. \quad (16)$$

From (14), (15) and (16), we obtain

$$\|u^1(\cdot, t) - u^2(\cdot, t)\| \leq 2^{(1-\frac{t}{T})} \delta^{\frac{t}{T}} E^{1-\frac{t}{T}}, \quad \forall t \in [0, T].$$

Hence, the theorem is proved. □

Next, we will take an example of specific functions $u^1(x, t)$, $u^2(x, t)$ satisfying the assumptions of Theorem 2.2.

Example 2.3. Let l, T, δ and E be the positive constants mentioned in (1) and in Theorem 2.2. Choose n_0 to be a positive integer satisfying

$$n_0 \geq \max \left\{ \frac{\sqrt{l}}{E}, \frac{l}{2\pi} \sqrt{\frac{1}{T} \ln \frac{l}{\delta^2}} \right\}. \quad (17)$$

Consider the functions

$$u^1(x, t) = \frac{1}{n_0} e^{-(\frac{2\pi n_0}{l})^2 t} \sin \frac{2\pi n_0 x}{l}, \quad 0 \leq x \leq l, 0 \leq t \leq T \quad (18)$$

$$u^2(x, t) = 0, \quad 0 \leq x \leq l, 0 \leq t \leq T. \quad (19)$$

It is easy to check that for $i = 1, 2$

$$\begin{cases} u_t^i - u_{xx}^i = 0, & 0 < x < l, 0 < t < T \\ u^i(0, t) = 0, & \int_0^l u^i(x, t) dx = 0, & 0 \leq t \leq T. \end{cases} \quad (20)$$

Furthermore

$$\begin{aligned} u_x^1(x, t) &= \frac{2\pi}{l} e^{-(\frac{2\pi n_0}{l})^2 t} \cos \frac{2\pi n_0 x}{l} \\ u_x^2(x, t) &= 0. \end{aligned}$$

This implies that

$$\begin{aligned} u_x^1(0, t) &= \frac{2\pi}{l} e^{-(\frac{2\pi n_0}{l})^2 t} \\ u_x^2(0, t) &= 0. \end{aligned}$$

It follows from $u^1(l, t) = u^2(l, t) = 0$ that

$$u^i(l, t) = C u_x^i(0, t), \quad i = 1, 2 \text{ with } C = 0.$$

Now we evaluate $\|u^i(\cdot, 0)\|, i = 1, 2$. It is clear that $\|u^2(\cdot, 0)\| = 0 < E$. We have

$$\begin{aligned} \|u^1(\cdot, 0)\|^2 &= \int_0^l (u^1(x, 0))^2 dx \\ &= \int_0^l \left(\frac{1}{n_0}\right)^2 \sin^2 \frac{2\pi n_0 x}{l} dx \\ &= \left(\frac{1}{n_0}\right)^2 \int_0^l \sin^2 \frac{2\pi n_0 x}{l} dx \\ &\leq \left(\frac{1}{n_0}\right)^2 \int_0^l dx = \frac{l}{n_0^2} \\ &\leq E^2 \text{ (since } n_0 \geq \frac{\sqrt{l}}{E} \text{)}. \end{aligned}$$

Therefore, we obtain $\|u^1(\cdot, 0)\| \leq E$. Next we evaluate $\|u^1(\cdot, T) - u^2(\cdot, T)\|$. We have

$$\begin{aligned} \|u^1(\cdot, T) - u^2(\cdot, T)\|^2 &= \int_0^l (u^1(x, T) - u^2(x, T))^2 dx \\ &= \int_0^l \frac{1}{n_0^2} e^{-(\frac{2\pi n_0}{l})^2 T} \sin^2 \frac{2\pi n_0 x}{l} dx \\ &\leq \frac{1}{n_0^2} e^{-(\frac{2\pi n_0}{l})^2 T} \int_0^l \sin^2 \frac{2\pi n_0 x}{l} dx \\ &\leq \frac{1}{n_0^2} e^{-(\frac{2\pi n_0}{l})^2 T} \int_0^l dx \\ &= \frac{1}{n_0^2} e^{-(\frac{2\pi n_0}{l})^2 T} l \\ &\leq e^{-(\frac{2\pi n_0}{l})^2 T} l \\ &\leq \delta^2 \text{ (since } n_0 \geq \frac{l}{2\pi} \sqrt{\frac{1}{T} \ln \frac{l}{\delta^2}} \text{)}. \end{aligned}$$

This implies that $\|u^1(\cdot, T) - u^2(\cdot, T)\| \leq \delta$.

Thus, the functions $u^1(x, t)$ and $u^2(x, t)$ satisfy the assumptions of Theorem 2.2. Therefore, we have

$$\|u^1(\cdot, t) - u^2(\cdot, t)\| \leq 2^{(1-\frac{t}{T})} \delta^{\frac{t}{T}} E^{1-\frac{t}{T}}, \forall t \in [0, T].$$

Acknowledgments: The work was supported by the Vietnam Ministry of Education and Training under Grant No. B2024-TDV-12.

REFERENCES

- [1] N. V. Duc, “Stability estimate for the heat equation backward in with Neumann and integral boundary conditions,” *Vinh University Journal of Science*, vol. 53, 3A, pp. 102–110, 2024. DOI: 10.56824/vujs.2024a055a
- [2] D. N. Hao, B. V. Huong, N. T. N. Oanh and P. X. Thanh, “Determination of a term in the right-hand side of parabolic equations,” *Journal of Computational and Applied Mathematics*, 309, pp. 28–43, 2017.
- [3] M. I. Ismailov, F. Kanca and D. Lesnic, “Determination of a time-dependent heat source under nonlocal boundary and integral overdetermination conditions,” *Applied Mathematics and Computation*, 218, pp. 4138–4146, 2011.
- [4] M. I. Ismailov, “Inverse Source Problem for Heat Equation with Nonlocal Wentzell Boundary Condition,” *Results Math*, 73(68), pp. 1422–6383, 2018.
- [5] F. Kanca and Mansur I. Ismailov, “The inverse problem of finding the time dependent diffusion coefficient of the heat equation from integral overdetermination data,” *Inverse Problems in Science and Engineering*, 20(4), pp. 463–476, 2012.
- [6] N. T. N. Oanh, “A method for choosing the regularization parameter of determining the right-hand side from integral observation,” *Asian-European Journal of Mathematics*, 15(7), 2250132, 2022.
- [7] N. T. N. Oanh and L. T. T. Giang, “Existence of optimal solution to the problem of finding an initial condition for nonlinear diffusion from integral observations,” *Asian-European Journal of Mathematics*, 14(8), 2150142, 2021.

TÓM TẮT

TÍNH ĐẶT KHÔNG CHÍNH VÀ ĐÁNH GIÁ ỔN ĐỊNH CHO PHƯƠNG TRÌNH TRUYỀN NHIỆT NGƯỢC THỜI GIAN VỚI CÁC ĐIỀU KIỆN BIÊN DIRICHLET VÀ TÍCH PHẦN

Nguyễn Văn Đức¹, Phan Hoài Linh²

¹ Khoa Toán, Trường Sư phạm, Trường Đại học Vinh, Việt Nam

² Cao học khoá 30 Toán Giải tích, Khoa Toán, Trường Đại Học Vinh, Nghệ An, Việt Nam

Ngày nhận bài 05/8/2024, ngày nhận đăng 17/10/2024

Trong bài báo này, đầu tiên chúng tôi chứng minh phương trình truyền nhiệt ngược thời gian với các điều kiện biên Dirichlet và tích phân là một bài toán đặt không chính. Sau đó, chúng tôi thành lập kết quả đánh giá ổn định kiểu Hölder cho bài toán này.

Từ khóa: Phương trình truyền nhiệt ngược; bài toán đặt không chính; đánh giá ổn định.